

CCDC73 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55304**Specification**

CCDC73 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O6ZRK6
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	124154

CCDC73 Polyclonal Antibody - Additional Information**Gene ID** 493860**Other Names**

Coiled-coil domain-containing protein 73, Sarcoma antigen NY-SAR-79, CCDC73

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CCDC73 Polyclonal Antibody - Protein Information**Name** CCDC73**CCDC73 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCDC73 Polyclonal Antibody - Images

Sample:U937(Human)Cell Lysate at 40 ug
Primary: Anti-CCDC73(bs-13812R)at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 124 kD
Observed band size: 124 kD



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CCDC73) Polyclonal Antibody, Unconjugated (bs-13812R) at 1:5000 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.