

CHST7 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55362

Specification

CHST7 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O9NS84
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54266

CHST7 Polyclonal Antibody - Additional Information

Gene ID 56548

Other Names

Carbohydrate sulfotransferase 7, 2.8.2.-, 2.8.2.17, Chondroitin 6-sulfotransferase 2, C6ST-2, Galactose/N-acetylglucosamine/N-acetylglucosamine 6-O-sulfotransferase 5, GST-5, N-acetylglucosamine 6-O-sulfotransferase 4, GlcNAc6ST-4, Gn6st-4, CHST7

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CHST7 Polyclonal Antibody - Protein Information

Name CHST7

Function

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of non-reducing N-acetylglucosamine (GlcNAc) residues. Preferentially acts on mannose-linked GlcNAc. Also able to catalyze the transfer of sulfate to position 6 of the N-acetylgalactosamine (GalNAc) residue of chondroitin. Also acts on core 2 mucin-type oligosaccharide and N-acetylglucosamine oligomer with a lower efficiency. Has weak or no activity toward keratan sulfate and oligosaccharides containing the Galbeta1-4GlcNAc. Catalyzes 6-O-sulfation of beta-benzyl GlcNAc but not alpha- or beta-benzyl GalNAc.

Cellular Location

Golgi apparatus membrane; Single-pass type II membrane protein

Tissue Location


Widely expressed. Highly expressed in heart, spleen, liver and ovary. Expressed at lower level in brain, placenta, thyroid, spinal cord and peripheral blood leukocytes. Not expressed in adult skin.

CHST7 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CHST7 Polyclonal Antibody - Images



Sample:

Kidney (Mouse) Lysate at 40 ug


Cerebrum (Mouse) Lysate at 40 ug

Primary: Anti- CHST7 (bs-13968R) at 1/1000 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 54 kD

Observed band size: 54 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CHST7) Polyclonal Antibody, Unconjugated (bs-13968R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

Ovary (Mouse) Lysate at 40 ug

Primary: Anti- CHST7 (bs-13968R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 54 kD

Observed band size: 54 kD