

CORO2B Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55374**Specification**

CORO2B Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9UQ03
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54953

CORO2B Polyclonal Antibody - Additional Information**Gene ID** 10391**Other Names**

Coronin-2B, Coronin-like protein C, Clipin-C, Protein FC96, CORO2B, KIAA0925

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CORO2B Polyclonal Antibody - Protein Information**Name** CORO2B**Synonyms** KIAA0925**Function**

May play a role in the reorganization of neuronal actin structure.

Cellular Location

Cytoplasm, cytoskeleton.

Tissue Location


Expressed predominantly in brain.

CORO2B Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CORO2B Polyclonal Antibody - Images



Tissue/cell: Mouse brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CORO2B Polyclonal Antibody, Unconjugated(bs-13997R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human gliomas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CORO2B Polyclonal Antibody, Unconjugated(bs-13997R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining