

ENOX1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55639

Specification

ENOX1 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q8TC92</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 73348

ENOX1 Polyclonal Antibody - Additional Information

Gene ID 55068

Other Names

Ecto-NOX disulfide-thiol exchanger 1, Candidate growth-related and time keeping constitutive hydroquinone [NADH] oxidase, cCNOX, Cell proliferation-inducing gene 38 protein, Constitutive Ecto-NOX, cNOX, Hydroquinone [NADH] oxidase, 1.-.-.-, Protein disulfide-thiol oxidoreductase, 1.-.-.-, ENOX1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

ENOX1 Polyclonal Antibody - Protein Information

Name ENOX1

Function

Probably acts as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 24 minutes and play a role in control of the ultradian cellular biological clock.

Cellular Location

Cell membrane. Secreted, extracellular space Note=Extracellular and plasma membrane-associated

Tissue Location

Expressed in lymphocyte cells, breast and breast cancer (at protein level). Found in the sera of cancer patients with a wide variety of cancers including breast, prostate, lung and ovarian



Tel: 858.875.1900 Fax: 858.875.1999

cancers, leukemias, and lymphomas. Found also in the serum of healthy volunteers or patients with disorders other than cancer. Probably shed into serum by cancer cells

ENOX1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ENOX1 Polyclonal Antibody - Images