

C8orf72 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55931

Specification

C8orf72 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q8TC76
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40728

C8orf72 Polyclonal Antibody - Additional Information

Gene ID 90362

Other Names

Protein FAM110B, FAM110B, C8orf72

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C8orf72 Polyclonal Antibody - Protein Information

Name FAM110B

Synonyms C8orf72

Function

May be involved in tumor progression.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome

Tissue Location


Detected in thyroid, spleen and testis, and at lower levels in stomach, spinal cord, lymph node, trachea, adrenal gland, prostate, ovary and intestine.

C8orf72 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C8orf72 Polyclonal Antibody - Images



Tissue/cell: human thyroid tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti- C8orf72 Polyclonal Antibody, Unconjugated(bs-15299R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining