

FBX40 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56091

Specification

FBX40 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9UH90
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79782

FBX40 Polyclonal Antibody - Additional Information

Gene ID 51725

Other Names

F-box only protein 40, Muscle disease-related protein, FBXO40, FBX40, KIAA1195

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FBX40 Polyclonal Antibody - Protein Information

Name FBXO40

Synonyms FBX40, KIAA1195

Function

Probable substrate-recognition component of the SCF (SKP1- CUL1-F-box protein)-type E3 ubiquitin ligase complex that may function in myogenesis.

Cellular Location

Cytoplasm.

Tissue Location


Expressed only in heart and skeletal muscle.

FBX40 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FBX40 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FBX40) Polyclonal Antibody, Unconjugated (bs-16049R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FBX40) Polyclonal Antibody, Unconjugated (bs-16049R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.