

GSTM3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56222**Specification**

GSTM3 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P21266
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26560

GSTM3 Polyclonal Antibody - Additional Information**Gene ID** 2947**Other Names**

Glutathione S-transferase Mu 3, 2.5.1.18, GST class-mu 3, GSTM3-3, hGSTM3-3, GSTM3, GST5

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

GSTM3 Polyclonal Antibody - Protein Information**Name** GSTM3**Synonyms** GST5**Function**

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. May govern uptake and detoxification of both endogenous compounds and xenobiotics at the testis and brain blood barriers.

Cellular Location

Cytoplasm.

Tissue Location


Testis and brain.

GSTM3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GSTM3 Polyclonal Antibody - Images



Tissue/cell: rat testis tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-GSTM3 Polyclonal Antibody, Unconjugated(bs-16341R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining