

KMT2B Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56406

Specification

KMT2B Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O9UMN6
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	293515

KMT2B Polyclonal Antibody - Additional Information

Gene ID 9757

Other Names

Histone-lysine N-methyltransferase 2B, Lysine N-methyltransferase 2B, 2.1.1.354, Myeloid/lymphoid or mixed-lineage leukemia protein 4, Trithorax homolog 2, WW domain-binding protein 7, WBP-7, KMT2B, HRX2, KIAA0304, MLL2, MLL4, TRX2, WBP7

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KMT2B Polyclonal Antibody - Protein Information

Name KMT2B

Synonyms HRX2, KIAA0304, MLL2, MLL4, TRX2, WBP7

Function

Histone methyltransferase that catalyzes methyl group transfer from S-adenosyl-L-methionine to the epsilon-amino group of 'Lys-4' of histone H3 (H3K4) via a non-processive mechanism. Part of chromatin remodeling machinery predominantly forms H3K4me1 and H3K4me2 methylation marks at active chromatin sites where transcription and DNA repair take place (PubMed: 17707229, PubMed: 25561738). Likely plays a redundant role with KMT2C in enriching H3K4me1 marks on primed and active enhancer elements (PubMed: 24081332). Plays a central role in beta-globin locus transcription regulation by being recruited by NFE2 (PubMed: 17707229). Plays an important role in controlling bulk H3K4me during oocyte growth and preimplantation development (By similarity). Required during the

transcriptionally active period of oocyte growth for the establishment and/or maintenance of bulk H3K4 trimethylation (H3K4me3), global transcriptional silencing that precedes resumption of meiosis, oocyte survival and normal zygotic genome activation (By similarity).

Cellular Location

Nucleus.

Tissue Location

Widely expressed. Highest levels in testis. Also found in brain with higher expression in the cerebellum than in any other region, bone marrow, heart, muscle, kidney, placenta, spleen, thymus, prostate, ovary, intestine, colon, peripheral blood lymphocytes and pancreas. Often amplified in pancreatic carcinomas

KMT2B Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KMT2B Polyclonal Antibody - Images