

SCGF Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56646**Specification**

SCGF Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9Y240
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35695

SCGF Polyclonal Antibody - Additional Information**Gene ID** 6320**Other Names**

C-type lectin domain family 11 member A, C-type lectin superfamily member 3, Lymphocyte secreted C-type lectin, Osteolectin, Stem cell growth factor, p47, CLEC11A, CLECSF3, LSLCL, SCGF

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SCGF Polyclonal Antibody - Protein Information**Name** CLEC11A**Synonyms** CLECSF3, LSLCL, SCGF**Function**

Promotes osteogenesis by stimulating the differentiation of mesenchymal progenitors into mature osteoblasts (PubMed:27976999). Important for repair and maintenance of adult bone (By similarity).

Cellular Location

Cytoplasm. Secreted

Tissue Location


Expressed in skeletal tissues including bone marrow, chondrocytes, primary ossification center-associated cells, the perichondrium and periosteum. Lower levels of expression were detected in spleen, thymus, appendix and fetal liver

SCGF Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SCGF Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SCGF) Polyclonal Antibody, Unconjugated (bs-17284R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.