

Histone H2A.V Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56697**Specification**

Histone H2A.V Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O71UI9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	13509

Histone H2A.V Polyclonal Antibody - Additional Information**Gene ID** 94239**Other Names**

Histone H2A.V, H2A.F/Z, H2A.Z variant histone 2 {ECO:0000312|HGNC:HGNC:20664}, H2AZ2 ([HGNC:20664](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=20664))

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Histone H2A.V Polyclonal Antibody - Protein Information**Name** H2AZ2 ([HGNC:20664](#))**Function**

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division (By similarity).

Cellular Location

Nucleus. Chromosome.

Histone H2A.V Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H2A.V Polyclonal Antibody - Images