

SPTBN3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56774

Specification

SPTBN3 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9H254
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	288985

SPTBN3 Polyclonal Antibody - Additional Information

Gene ID 57731

Other Names

Spectrin beta chain, non-erythrocytic 4, Beta-IV spectrin, Spectrin, non-erythroid beta chain 3, SPTBN4, KIAA1642, SPTBN3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SPTBN3 Polyclonal Antibody - Protein Information

Name SPTBN4

Synonyms KIAA1642, SPTBN3

Cellular Location

Cytoplasm, cytoskeleton. Cytoplasm, cell cortex

Tissue Location

Expressed in skeletal muscle at the sarcolemma and in the muscle capillaries (at protein level) (PubMed:28540413) Abundantly expressed in brain and pancreatic islets (PubMed:11086001)


SPTBN3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPTBN3 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SPTBN3) Polyclonal Antibody, Unconjugated (bs-17649R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.