

MOXD1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56811

Specification

MOXD1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q6UVY6
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	69652

MOXD1 Polyclonal Antibody - Additional Information

Gene ID 26002

Other Names

DBH-like monooxygenase protein 1, 1.14.17.-, Monooxygenase X, MOXD1, MOX

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MOXD1 Polyclonal Antibody - Protein Information

Name MOXD1

Synonyms MOX

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein

Tissue Location

Highly expressed in lung, kidney, brain and spinal cord.


MOXD1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MOXD1 Polyclonal Antibody - Images



Sample:


Jurkat(Human) Cell Lysate at 30 ug

Primary: Anti-MOXD1 (bs-17733R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 68 kD

Observed band size: 68 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MOXD1) Polyclonal Antibody, Unconjugated (bs-17733R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.