

ZNF622 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57123

Specification

ZNF622 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O969S3
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54272

ZNF622 Polyclonal Antibody - Additional Information

Gene ID 90441

Other Names

Zinc finger protein 622, Zinc finger-like protein 9, ZNF622, ZPR9

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZNF622 Polyclonal Antibody - Protein Information

Name ZNF622 {ECO:0000303|PubMed:32669547, ECO:0000312|HGNC:HGNC:30958}

Function

Pre-60S-associated cytoplasmic factor involved in the cytoplasmic maturation of the 60S subunit.

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Expressed in lung, kidney, spleen, liver and brain with lowest expression in kidney.


ZNF622 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF622 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF622) Polyclonal Antibody, Unconjugated (bs-18514R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.