

**ZCCHV Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57151****Specification**

---

**ZCCHV Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q7Z2W4</a>
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	101431

**ZCCHV Polyclonal Antibody - Additional Information****Gene ID** 56829**Other Names**

Zinc finger CCCH-type antiviral protein 1, ADP-ribosyltransferase diphtheria toxin-like 13, ARTD13, Inactive Poly [ADP-ribose] polymerase 13, PARP13, Zinc finger CCCH domain-containing protein 2, Zinc finger antiviral protein, ZAP, ZC3HAV1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=23721](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=23721)), ZC3HDC2

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ZCCHV Polyclonal Antibody - Protein Information****Name** ZC3HAV1 ([HGNC:23721](#))**Synonyms** ZC3HDC2**Function**

Antiviral protein which inhibits the replication of viruses by recruiting the cellular RNA degradation machineries to degrade the viral mRNAs. Binds to a ZAP-responsive element (ZRE) present in the target viral mRNA, recruits cellular poly(A)-specific ribonuclease PARN to remove the poly(A) tail, and the 3'-5' exoribonuclease complex exosome to degrade the RNA body from the 3'-end. It also recruits the decapping complex DCP1-DCP2 through RNA helicase p72 (DDX17) to remove the cap structure of the viral mRNA to initiate its degradation from the 5'-end. Its target viruses belong to families which include retroviridae: human immunodeficiency virus type 1 (HIV-1), moloney and murine leukemia virus (MoMLV) and xenotropic MuLV-related virus (XMRV), filoviridae: ebola virus (EBOV) and marburg virus (MARV), togaviridae: sindbis virus (SINV) and Ross river virus (RRV). Specifically targets the multiply spliced but not unspliced or singly spliced HIV-1 mRNAs for

degradation. Isoform 1 is a more potent viral inhibitor than isoform 2. Isoform 2 acts as a positive regulator of RIGI signaling resulting in activation of the downstream effector IRF3 leading to the expression of type I IFNs and IFN stimulated genes (ISGs).

**Cellular Location**

[Isoform 1]: Cytoplasm {ECO:0000250|UniProtKB:Q8K3Y6}. Nucleus

{ECO:0000250|UniProtKB:Q8K3Y6} Note=Localizes in the cytoplasm at steady state, but shuttles between nucleus and cytoplasm in a XPO1-dependent manner {ECO:0000250|UniProtKB:Q8K3Y6}

**ZCCHV Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ZCCHV Polyclonal Antibody - Images**