

MED12L Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57253

Specification

MED12L Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E
Primary Accession Q86YW9
Host Rabbit
Clonality Polyclonal
Calculated MW 240120

MED12L Polyclonal Antibody - Additional Information

Gene ID 116931

Other Names

Mediator of RNA polymerase II transcription subunit 12-like protein, Mediator complex subunit 12-like protein, Thyroid hormone receptor-associated-like protein, Trinucleotide repeat-containing gene 11 protein-like, MED12L, KIAA1635, TNRC11L, TRALP, TRALPUSH

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

MED12L Polyclonal Antibody - Protein Information

Name MED12L

Synonyms KIAA1635, TNRC11L, TRALP, TRALPUSH


Function

May be a component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity).

Cellular Location

Nucleus.

MED12L Polyclonal Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MED12L Polyclonal Antibody - Images