

**LCAT Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57642****Specification**

---

**LCAT Polyclonal Antibody - Product Information**

|                   |                              |
|-------------------|------------------------------|
| Application       | <b>WB, IHC-P</b>             |
| Primary Accession | <a href="#">P04180</a>       |
| Reactivity        | <b>Rat, Pig, Dog, Bovine</b> |
| Host              | <b>Rabbit</b>                |
| Clonality         | <b>Polyclonal</b>            |
| Calculated MW     | <b>49578</b>                 |

**LCAT Polyclonal Antibody - Additional Information****Gene ID** 3931**Other Names**

Phosphatidylcholine-sterol acyltransferase, 2.3.1.43, 1-alkyl-2-acetylgllycerophosphocholine esterase, 3.1.1.47, Lecithin-cholesterol acyltransferase, Phospholipid-cholesterol acyltransferase, Platelet-activating factor acetylhydrolase, PAF acetylhydrolase, LCAT

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**LCAT Polyclonal Antibody - Protein Information****Name** LCAT**Function**

Central enzyme in the extracellular metabolism of plasma lipoproteins. Synthesized mainly in the liver and secreted into plasma where it converts cholesterol and phosphatidylcholines (lecithins) to cholesteryl esters and lysophosphatidylcholines on the surface of high and low density lipoproteins (HDLs and LDLs) (PubMed: [10329423](http://www.uniprot.org/citations/10329423), PubMed: [19065001](http://www.uniprot.org/citations/19065001), PubMed: [26195816](http://www.uniprot.org/citations/26195816)). The cholesterol ester is then transported back to the liver. Has a preference for plasma 16:0-18:2 or 18:0-18:2 phosphatidylcholines (PubMed: [8820107](http://www.uniprot.org/citations/8820107)). Also produced in the brain by primary astrocytes, and esterifies free cholesterol on nascent APOE-containing lipoproteins secreted from glia and influences cerebral spinal fluid (CSF) APOE- and APOA1 levels. Together with APOE and the cholesterol transporter ABCA1, plays a key role in the maturation of glial-derived, nascent lipoproteins. Required for remodeling high- density lipoprotein particles into their spherical forms (PubMed: [10722751](http://www.uniprot.org/citations/10722751))

target="\_blank">10722751</a>). Catalyzes the hydrolysis of 1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine (platelet-activating factor or PAF) to 1-O-alkyl-sn-glycero-3-phosphocholine (lyso-PAF) (PubMed:<a href="http://www.uniprot.org/citations/8016111" target="\_blank">8016111</a>). Also catalyzes the transfer of the acetate group from PAF to 1-hexadecanoyl-sn-glycero-3-phosphocholine forming lyso-PAF (PubMed:<a href="http://www.uniprot.org/citations/8016111" target="\_blank">8016111</a>). Catalyzes the esterification of (24S)-hydroxycholesterol (24(S)OH-C), also known as cerebrosterol to produce 24(S)OH-C monoesters (PubMed:<a href="http://www.uniprot.org/citations/24620755" target="\_blank">24620755</a>).

#### Cellular Location

Secreted. Note=Secreted into blood plasma (PubMed:10222237, PubMed:3458198, PubMed:8820107) Produced in astrocytes and secreted into cerebral spinal fluid (CSF) (PubMed:10222237).

#### Tissue Location


Detected in blood plasma (PubMed:10222237, PubMed:3458198, PubMed:8820107). Detected in cerebral spinal fluid (at protein level) (PubMed:10222237). Detected in liver (PubMed:3458198, PubMed:3797244). Expressed mainly in brain, liver and testes

### LCAT Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### LCAT Polyclonal Antibody - Images




Sample:  
Liver (Mouse) Lysate at 40 ug


Primary: Anti- LCAT (bs-1972R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 47 kD  
Observed band size: 49kD




Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;  
Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;  
Incubation: Anti-LCAT Polyclonal Antibody, Unconjugated(bs-1972R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Sample: K562 (human)Cell Lysate at 40 ug  
Primary: Anti- LCAT(bs-1972R)at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 47kD  
Observed band size: 47/52 kD



Sample: brain (mouse) Lysate at 40 ug  
 Primary: Anti- LCAT(bs-1972R)at 1/300 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
 Predicted band size: 47kD  
 Observed band size: 48kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LCAT) Polyclonal Antibody, Unconjugated (bs-1972R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20

minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LCAT) Polyclonal Antibody, Unconjugated (bs-1972R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.