

PABPC4 Polyclonal Antibody Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57747

Specification

PABPC4 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC-P, IHC-F, IF, ICC, E <u>013310</u> Rat, Pig, Dog, Bovine Rabbit Polyclonal 70783

PABPC4 Polyclonal Antibody - Additional Information

Gene ID 8761

Other Names Polyadenylate-binding protein 4, PABP-4, Poly(A)-binding protein 4, Activated-platelet protein 1, APP-1, Inducible poly(A)-binding protein, iPABP, PABPC4, APP1, PABP4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PABPC4 Polyclonal Antibody - Protein Information

Name PABPC4

Synonyms APP1, PABP4

Function

Binds the poly(A) tail of mRNA (PubMed:8524242). Binds to SMIM26 mRNA and plays a role in its post-transcriptional regulation (PubMed:37009826). May be involved in cytoplasmic regulatory processes of mRNA metabolism. Can probably bind to cytoplasmic RNA sequences other than poly(A) in vivo (By similarity).

Cellular Location

Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs.

Tissue Location

Expressed at low levels in resting normal T cells; following T-cell activation, however, mRNA levels are rapidly up- regulated



PABPC4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PABPC4 Polyclonal Antibody - Images