

DLC2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57767

Specification

DLC2 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	O9Y3M8
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	124967

DLC2 Polyclonal Antibody - Additional Information

Gene ID 90627

Other Names

StAR-related lipid transfer protein 13, 46H23.2, Deleted in liver cancer 2 protein, DLC-2, Rho GTPase-activating protein, START domain-containing protein 13, StARD13, STARD13, DLC2, GT650

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DLC2 Polyclonal Antibody - Protein Information

Name STARD13

Synonyms DLC2, GT650

Function

GTPase-activating protein for RhoA, and perhaps for Cdc42. May be involved in regulation of cytoskeletal reorganization, cell proliferation and cell motility. Acts a tumor suppressor in hepatocellular carcinoma cells.

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein; Cytoplasmic side. Mitochondrion membrane; Peripheral membrane protein; Cytoplasmic side. Lipid droplet

Tissue Location

Ubiquitously expressed. Underexpressed in hepatocellular carcinoma cells and some breast cancer cell lines

DLC2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DLC2 Polyclonal Antibody - Images