

GRIM19 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58077**Specification**

GRIM19 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O9P0J0
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16698

GRIM19 Polyclonal Antibody - Additional Information**Gene ID** 51079**Other Names**

NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13, Cell death regulatory protein GRIM-19, Complex I-B16.6, CI-B16.6, Gene associated with retinoic and interferon-induced mortality 19 protein, GRIM-19, Gene associated with retinoic and IFN-induced mortality 19 protein, NADH-ubiquinone oxidoreductase B16.6 subunit, NDUFA13, GRIM19

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

GRIM19 Polyclonal Antibody - Protein Information**Name** NDUFA13**Synonyms** GRIM19**Function**

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis (PubMed:27626371). Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (PubMed:27626371). Involved in the interferon/all-trans-retinoic acid (IFN/RA) induced cell death. This apoptotic activity is inhibited by interaction with viral IRF1. Prevents the transactivation of STAT3 target genes. May play a role in CARD15-mediated innate mucosal responses and serve to regulate intestinal epithelial cell responses to microbes (PubMed:15753091).

Cellular Location

Mitochondrion inner membrane; Single-pass membrane protein; Matrix side. Nucleus
Note=Localizes mainly in the mitochondrion (PubMed:12628925). May be translocated into the nucleus upon IFN/RA treatment


Tissue Location

Widely expressed, with highest expression in heart, skeletal muscle, liver, kidney and placenta. In intestinal mucosa, down-regulated in areas involved in Crohn disease and ulcerative colitis.


GRIM19 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


GRIM19 Polyclonal Antibody - Images

Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GRIM19) Polyclonal Antibody, Unconjugated (bs-3594R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse heart tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide

for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GRIM19) Polyclonal Antibody, Unconjugated (bs-3594R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

Liver (Mouse) Lysate at 40 ug

Primary: Anti-GRIM19 (bs-3594R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 16 kD

Observed band size: 17 kD