

MEG1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58081

Specification

MEG1 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P29074
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105911

MEG1 Polyclonal Antibody - Additional Information

Gene ID 5775

Other Names

Tyrosine-protein phosphatase non-receptor type 4, 3.1.3.48, Protein-tyrosine phosphatase MEG1, MEG, PTPase-MEG1, PTPN4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MEG1 Polyclonal Antibody - Protein Information

Name PTPN4

Function

Phosphatase that plays a role in immunity, learning, synaptic plasticity or cell homeostasis (PubMed:25825441, PubMed:27246854). Regulates neuronal cell homeostasis by protecting neurons against apoptosis (PubMed:20086240). Negatively regulates TLR4-induced interferon beta production by dephosphorylating adapter TICAM2 and inhibiting subsequent TRAM-TRIF interaction (PubMed:25825441). Dephosphorylates also the immunoreceptor tyrosine-based activation motifs/ITAMs of the TCR zeta subunit and thereby negatively regulates TCR-mediated signaling pathway (By similarity). May act at junctions between the membrane and the cytoskeleton.

Cellular Location


Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton
Cytoplasm

MEG1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MEG1 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MEG1) Polyclonal Antibody, Unconjugated (bs-3629R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.