

NDUFS1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58106

Specification

NDUFS1 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P28331
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	79468

NDUFS1 Polyclonal Antibody - Additional Information

Gene ID 4719

Other Names

NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial, 7.1.1.2, Complex I-75kD, CI-75kD, NDUFS1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NDUFS1 Polyclonal Antibody - Protein Information

Name NDUFS1

Function

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed: [30879903](http://www.uniprot.org/citations/30879903), PubMed: [31557978](http://www.uniprot.org/citations/31557978)). Essential for catalysing the entry and efficient transfer of electrons within complex I (PubMed: [31557978](http://www.uniprot.org/citations/31557978)). Plays a key role in the assembly and stability of complex I and participates in the association of complex I with ubiquinol-cytochrome reductase complex (Complex III) to form supercomplexes (PubMed: [30879903](http://www.uniprot.org/citations/30879903), PubMed: [31557978](http://www.uniprot.org/citations/31557978)).

Cellular Location


Mitochondrion inner membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P15690}; Matrix side {ECO:0000250|UniProtKB:P15690}

NDUFS1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDUFS1 Polyclonal Antibody - Images



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-NDUFS1/NADH dehydrogenase (ubiquinone) FeS protein 1(75kD) Polyclonal Antibody, Unconjugated(bs-3890R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining