

**FBP2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58114****Specification**

---

**FBP2 Polyclonal Antibody - Product Information**

Application	<b>IHC-P</b>
Primary Accession	<a href="#">O00757</a>
Reactivity	<b>Rat, Dog, Bovine</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>36743</b>

**FBP2 Polyclonal Antibody - Additional Information****Gene ID** 8789**Other Names**

Fructose-1, 6-bisphosphatase isozyme 2, FBPase 2, 3.1.3.11, D-fructose-1, 6-bisphosphate 1-phosphohydrolase 2, Muscle FBPase, FBP2

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**FBP2 Polyclonal Antibody - Protein Information****Name** FBP2**Function**

Catalyzes the hydrolysis of fructose 1,6-bisphosphate to fructose 6-phosphate in the presence of divalent cations and probably participates in glycogen synthesis from carbohydrate precursors, such as lactate.

**Cellular Location**

Cell junction. Cytoplasm. Nucleus. Cytoplasm, myofibril, sarcomere, Z line. Note=In neonatal cardiomyocytes, distributed throughout the cytosol, accumulating in the intercalated disks which occur at the Z line of cardiomyocytes and connect adjacent cells, and also located in the nucleus; dissociates from the Z line following an increase in cytosolic Ca(2+) concentration (By similarity). In muscle precursor cells, localizes predominantly to the nucleus and to a lesser extent to the cytoplasm at the proliferative phase, while mainly localizing to the cytoplasm at the differentiation phase (By similarity). Colocalizes with ALDOA and alpha-actinin on both sides of the Z line of skeletal muscle; dissociates rapidly from the Z line following an increase in cytosolic Ca(2+) concentration.

### Tissue Location


Expressed in skeletal muscle (at protein level).

### FBP2 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### FBP2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FBP2) Polyclonal Antibody, Unconjugated (bs-3981R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FBP2) Polyclonal Antibody, Unconjugated (bs-3981R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.