

ARHGAP20 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58143

Specification

ARHGAP20 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9P2F6
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	132608

ARHGAP20 Polyclonal Antibody - Additional Information

Gene ID 57569

Other Names

Rho GTPase-activating protein 20, Rho-type GTPase-activating protein 20, ARHGAP20, KIAA1391

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ARHGAP20 Polyclonal Antibody - Protein Information

Name ARHGAP20

Synonyms KIAA1391

Function

GTPase activator for the Rho-type GTPases by converting them to an inactive GDP-bound state.

Tissue Location

Expressed predominantly in the brain. Lower expression is found in lymph nodes.


ARHGAP20 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ARHGAP20 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ARHGAP20) Polyclonal Antibody, Unconjugated (bs-4271R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.