

**ACOX2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58264**

**Specification**

---

**ACOX2 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">O99424</a>
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76827

**ACOX2 Polyclonal Antibody - Additional Information**

Gene ID 8309

**Other Names**

Peroxisomal acyl-coenzyme A oxidase 2, 1.17.99.3, 3-alpha, 7-alpha, 12-alpha-trihydroxy-5-beta-cholestanoyl-CoA 24-hydroxylase, 3-alpha, 7-alpha, 12-alpha-trihydroxy-5-beta-cholestanoyl-CoA oxidase, THCCox, ACOX2 ([HGNC:120](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=120))

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ACOX2 Polyclonal Antibody - Protein Information**

Name ACOX2 ([HGNC:120](#))

**Function**

Oxidizes the CoA esters of the bile acid intermediates di- and tri-hydroxycholestanic acids (PubMed:[27884763](http://www.uniprot.org/citations/27884763)). Capable of oxidizing short as well as long chain 2-methyl branched fatty acids (By similarity).

**Cellular Location**

Peroxisome

**Tissue Location**


Present in all tissues tested: heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Most abundant in heart, liver and kidney.

## ACOX2 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## ACOX2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ACOX2) Polyclonal Antibody, Unconjugated (bs-5030R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ACOX2) Polyclonal Antibody, Unconjugated (bs-5030R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ACOX2) Polyclonal Antibody, Unconjugated (bs-5030R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.