

FADS1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58269

Specification

FADS1 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O60427
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51964

FADS1 Polyclonal Antibody - Additional Information

Gene ID 3992

Other Names

Acyl-CoA (8-3)-desaturase, 1.14.19.44, Delta(5) fatty acid desaturase, D5D, Delta-5 desaturase, Fatty acid desaturase 1, FADS1 {ECO:0000303|PubMed:10860662, ECO:0000312|HGNC:HGNC:3574}

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FADS1 Polyclonal Antibody - Protein Information

Name FADS1 {ECO:0000303|PubMed:10860662, ECO:0000312|HGNC:HGNC:3574}

Function

[Isoform 1]: Acts as a front-end fatty acyl-coenzyme A (CoA) desaturase that introduces a cis double bond at carbon 5 located between a preexisting double bond and the carboxyl end of the fatty acyl chain. Involved in biosynthesis of highly unsaturated fatty acids (HUFA) from the essential polyunsaturated fatty acids (PUFA) linoleic acid (LA) (18:2n-6) and alpha-linolenic acid (ALA) (18:3n-3) precursors. Specifically, desaturates dihomo-gamma-linoleoate (DGLA) (20:3n-6) and eicosatetraenoate (ETA) (20:4n-3) to generate arachidonate (AA) (20:4n-6) and eicosapentaenoate (EPA) (20:5n-3), respectively (PubMed:10601301, PubMed:10769175). As a rate limiting enzyme for DGLA (20:3n-6) and AA (20:4n-6)-derived eicosanoid biosynthesis, controls the metabolism of inflammatory lipids like prostaglandin E2, critical for efficient acute inflammatory response and maintenance of epithelium homeostasis. Contributes to membrane phospholipid biosynthesis by providing AA (20:4n-6) as a major acyl chain esterified into phospholipids. In particular, regulates phosphatidylinositol-4,5-bisphosphate levels, modulating inflammatory

cytokine production in T-cells (By similarity). Also desaturates (11E)- octadecenoate (trans-vaccenoate)(18:1n-9), a metabolite in the biohydrogenation pathway of LA (18:2n-6) (By similarity).

Cellular Location

[Isoform 1]: Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:A4UVI1}; Multi-pass membrane protein {ECO:0000250|UniProtKB:A4UVI1}. Mitochondrion

Tissue Location


Widely expressed, with highest levels in liver, brain, adrenal gland and heart. Highly expressed in fetal liver and brain.

FADS1 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FADS1 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FADS1) Polyclonal Antibody, Unconjugated (bs-5060R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

**Sample:**


Adrenal gland (Mouse) Lysate at 40 ug

Primary: Anti- FADS1 (bs-5060R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 49 kD

Observed band size: 50 kD

**Sample:**

Liver (Mouse) Lysate at 40 ug

Primary: Anti- FADS1 (bs-5060R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 49 kD

Observed band size: 50 kD