

**ARHGAP24 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58313**

### Specification

---

#### ARHGAP24 Polyclonal Antibody - Product Information

Application	WB, E
Primary Accession	<a href="#">Q8N264</a>
Reactivity	Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84258

#### ARHGAP24 Polyclonal Antibody - Additional Information

Gene ID 83478

#### Other Names

Rho GTPase-activating protein 24, Filamin-A-associated RhoGAP, FILGAP, RAC1- and CDC42-specific GTPase-activating protein of 72 kDa, RC-GAP72, Rho-type GTPase-activating protein 24, RhoGAP of 73 kDa, Sarcoma antigen NY-SAR-88, p73RhoGAP, ARHGAP24, FILGAP

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

#### ARHGAP24 Polyclonal Antibody - Protein Information

Name ARHGAP24

Synonyms FILGAP

#### Function

Rho GTPase-activating protein involved in cell polarity, cell morphology and cytoskeletal organization. Acts as a GTPase activator for the Rac-type GTPase by converting it to an inactive GDP-bound state. Controls actin remodeling by inactivating Rac downstream of Rho leading to suppress leading edge protrusion and promotes cell retraction to achieve cellular polarity. Able to suppress RAC1 and CDC42 activity in vitro. Overexpression induces cell rounding with partial or complete disruption of actin stress fibers and formation of membrane ruffles, lamellipodia, and filopodia. Isoform 2 is a vascular cell-specific GAP involved in modulation of angiogenesis.

#### Cellular Location

Cytoplasm, cytoskeleton. Cell junction, adherens junction. Cell junction, focal adhesion. Cell projection Note=Localizes to actin stress fibers. In migrating cells, localizes to membrane lamellae and protrusions

**Tissue Location**

Isoform 1 is widely expressed with a higher level in kidney. Isoform 2 is mainly expressed in endothelial cells

**ARHGAP24 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ARHGAP24 Polyclonal Antibody - Images**