

Ubiquitously expressed transcript Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58504

Specification

Ubiquitously expressed transcript Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9UBK9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18246

Ubiquitously expressed transcript Polyclonal Antibody - Additional Information

Gene ID 8409

Other Names

Protein UXT, Androgen receptor trapped clone 27 protein, ART-27, Ubiquitously expressed transcript protein, UXT

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Ubiquitously expressed transcript Polyclonal Antibody - Protein Information

Name UXT

Function

Involved in gene transcription regulation (PubMed:21730289, PubMed:28106301). Acts in concert with the corepressor URI1 to regulate androgen receptor AR-mediated transcription (PubMed:11854421, PubMed:21730289). Together with URI1, associates with chromatin to the NKX3-1 promoter region (PubMed:21730289). Negatively regulates the transcriptional activity of the estrogen receptor ESR1 by inducing its translocation into the cytoplasm (PubMed:28106301). May act as nuclear chaperone that facilitates the formation of the NF-kappa-B enhanceosome and thus positively regulates NF-kappa-B transcription activity (PubMed:17620405, PubMed:21307340). Potential component of mitochondrial-associated LRPPRC, a multidomain organizer that potentially

integrates mitochondria and the microtubular cytoskeleton with chromosome remodeling (PubMed:17554592). Increasing concentrations of UXT contributes to progressive aggregation of mitochondria and cell death potentially through its association with LRPPRC (PubMed:17554592). Suppresses cell transformation and it might mediate this function by interaction and inhibition of the biological activity of cell proliferation and survival stimulatory factors like MECOM (PubMed:17635584).

Cellular Location

[Isoform 1]: Cytoplasm Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole Note=Predominantly localizes to the nucleus (PubMed:16221885) Localizes to spindle pole during mitosis (PubMed:16221885)

Tissue Location


Ubiquitous (PubMed:10087202, PubMed:11827465, PubMed:11854421, PubMed:17635584). Expressed in prostate epithelial cells (PubMed:21730289). Expressed in mammary epithelial cells (PubMed:28106301). Highest levels in the heart, skeletal muscle, pancreas, kidney, liver, adrenal gland, peripheral blood leukocytes, lymph node, prostate, and thyroid and the lowest levels in bladder and uterus (PubMed:11827465, PubMed:11854421, PubMed:17635584) Overexpressed in a number of tumor tissues (PubMed:11854421, PubMed:16221885, PubMed:28106301).

Ubiquitously expressed transcript Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ubiquitously expressed transcript Polyclonal Antibody - Images




Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen


peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Protein UXT) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (Mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.