

TANK Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5860c

Specification

TANK Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O92844
Other Accession	NP_004171.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	47816
Antigen Region	88-114

TANK Antibody (Center) - Additional Information

Gene ID 10010

Other Names

TRAF family member-associated NF-kappa-B activator, TRAF-interacting protein, I-TRAF, TANK, ITRAF, TRAF2

Target/Specificity

This TANK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-114 amino acids from the Central region of human TANK.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TANK Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

TANK Antibody (Center) - Protein Information

Name TANK

Synonyms ITRAF, TRAF2

Function Adapter protein involved in I-kappa-B-kinase (IKK) regulation which constitutively binds TBK1 and IKBE playing a role in antiviral innate immunity. Acts as a regulator of TRAF function by maintaining them in a latent state. Blocks TRAF2 binding to LMP1 and inhibits LMP1- mediated NF-kappa-B activation. Negatively regulates NF-kappaB signaling and cell survival upon DNA damage (PubMed:[25861989](#)). Plays a role as an adapter to assemble ZC3H12A, USP10 in a deubiquitination complex which plays a negative feedback response to attenuate NF-kappaB activation through the deubiquitination of IKKKG or TRAF6 in response to interleukin-1-beta (IL1B) stimulation or upon DNA damage (PubMed:[25861989](#)). Promotes UBP10-induced deubiquitination of TRAF6 in response to DNA damage (PubMed:[25861989](#)). May control negatively TRAF2-mediated NF-kappa-B activation signaled by CD40, TNFR1 and TNFR2.

Cellular Location

Cytoplasm.

Tissue Location


Ubiquitous.

TANK Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TANK Antibody (Center) - Images



TANK Antibody (Center) (Cat. #AP5860c) western blot analysis in WiDr cell line lysates (35ug/lane). This demonstrates the TANK antibody detected the TANK protein (arrow).



TANK Antibody (Center) (Cat. #AP5860c) immunohistochemistry analysis in formalin fixed and paraffin embedded human cervix carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TANK Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



TANK Antibody (Center) (Cat. #AP5860c) flow cytometric analysis of WiDr cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.