

**ATG14 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58664**

**Specification**

---

**ATG14 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">O6ZNE5</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55309

**ATG14 Polyclonal Antibody - Additional Information**

Gene ID 22863

**Other Names**

Beclin 1-associated autophagy-related key regulator, Barkor, Autophagy-related protein 14-like protein, Atg14L, ATG14 {ECO:0000303|PubMed:18843052}

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ATG14 Polyclonal Antibody - Protein Information**

Name ATG14 {ECO:0000303|PubMed:18843052}

**Function**

Required for both basal and inducible autophagy. Determines the localization of the autophagy-specific PI3-kinase complex PI3KC3-C1 (PubMed: [18843052](http://www.uniprot.org/citations/18843052)), PubMed: [19050071](http://www.uniprot.org/citations/19050071)). Plays a role in autophagosome formation and MAP1LC3/LC3 conjugation to phosphatidylethanolamine (PubMed: [19270696](http://www.uniprot.org/citations/19270696)), PubMed: [20713597](http://www.uniprot.org/citations/20713597)). Promotes BECN1 translocation from the trans-Golgi network to autophagosomes (PubMed: [20713597](http://www.uniprot.org/citations/20713597)). Enhances PIK3C3 activity in a BECN1-dependent manner. Essential for the autophagy-dependent phosphorylation of BECN1 (PubMed: [23878393](http://www.uniprot.org/citations/23878393)). Stimulates the phosphorylation of BECN1, but suppresses the phosphorylation PIK3C3 by AMPK (PubMed: [23878393](http://www.uniprot.org/citations/23878393)). Binds to STX17-SNAP29 binary t-SNARE complex on autophagosomes and primes it for VAMP8 interaction to promote autophagosome-endolysosome

fusion (PubMed:<a href="http://www.uniprot.org/citations/25686604" target="\_blank">25686604</a>, PubMed:<a href="http://www.uniprot.org/citations/37632749" target="\_blank">37632749</a>). Modulates the hepatic lipid metabolism (By similarity).

#### **Cellular Location**

Cytoplasm. Endoplasmic reticulum membrane; Peripheral membrane protein. Preautophagosomal structure membrane; Peripheral membrane protein. Cytoplasmic vesicle, autophagosome membrane; Peripheral membrane protein. Note=Cytosolic under nutrient-rich conditions (PubMed:19050071). Following autophagy stimuli, such as starvation or rapamycin induction, predominantly detected in cytoplasmic foci, identified as isolation membranes and autophagosomes (PubMed:19050071). Accumulates on highly curved PtdIns(3)P enriched autophagic membrane via its BATS domain to sense and maintain membrane curvature (By similarity). Localizes also to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme (By similarity). {ECO:0000250|UniProtKB:Q8CDJ3}

#### **ATG14 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ATG14 Polyclonal Antibody - Images**