

ZHX2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58684

Specification

ZHX2 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9Y6X8
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92307

ZHX2 Polyclonal Antibody - Additional Information

Gene ID 22882

Other Names

Zinc fingers and homeoboxes protein 2, Alpha-fetoprotein regulator 1, AFP regulator 1, Regulator of AFP, Zinc finger and homeodomain protein 2, ZHX2, AFR1, KIAA0854, RAF

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZHX2 Polyclonal Antibody - Protein Information

Name ZHX2

Synonyms AFR1, KIAA0854, RAF

Function

Acts as a transcriptional repressor (PubMed: [12741956](http://www.uniprot.org/citations/12741956)). Represses the promoter activity of the CDC25C gene stimulated by NFYA (PubMed: [12741956](http://www.uniprot.org/citations/12741956)). May play a role in retinal development where it regulates the composition of bipolar cell populations, by promoting differentiation of bipolar OFF-type cells (By similarity). In the brain, may promote maintenance and suppress differentiation of neural progenitor cells in the developing cortex (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:12741956, ECO:0000269|PubMed:17056598} Note=Colocalizes with EFNB1 intracellular domain in the nucleus {ECO:0000250|UniProtKB:Q8C0C0}


Tissue Location

Ubiquitously expressed. Expressed in podocytes.


ZHX2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZHX2 Polyclonal Antibody - Images

Paraformaldehyde-fixed, paraffin embedded (Rat thyroid); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZHX2) Polyclonal Antibody, Unconjugated (bs-7581R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZHX2) Polyclonal Antibody, Unconjugated (bs-7581R) at 1:400 overnight at 4°C, followed by operating

according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.