

**Blood Group Antigen Precursor Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
Catalog # AP58686

**Specification**

---

**Blood Group Antigen Precursor Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">P16442</a>
Reactivity	Chimpanzee
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40934

**Blood Group Antigen Precursor Polyclonal Antibody - Additional Information**

Gene ID 28

**Other Names**

Histo-blood group ABO system transferase, Fucosylglycoprotein 3-alpha-galactosyltransferase, Fucosylglycoprotein alpha-N-acetylgalactosaminyltransferase, Glycoprotein-fucosylgalactoside alpha-N-acetylgalactosaminyltransferase, 2.4.1.40, Glycoprotein-fucosylgalactoside alpha-galactosyltransferase, 2.4.1.37, Histo-blood group A transferase, A transferase, Histo-blood group B transferase, B transferase, NAGAT, Fucosylglycoprotein alpha-N-acetylgalactosaminyltransferase soluble form, ABO

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Blood Group Antigen Precursor Polyclonal Antibody - Protein Information**

Name ABO

**Function**

This protein is the basis of the ABO blood group system. The histo-blood group ABO involves three carbohydrate antigens: A, B, and H. A, B, and AB individuals express a glycosyltransferase activity that converts the H antigen to the A antigen (by addition of UDP-GalNAc) or to the B antigen (by addition of UDP-Gal), whereas O individuals lack such activity.

**Cellular Location**

Golgi apparatus, Golgi stack membrane; Single- pass type II membrane protein. Secreted Note=Membrane-bound form in trans cisternae of Golgi. Secreted into the body fluid

**Tissue Location**

Expressed at high levels in testis. Also expressed in pancreas, uterus and lung and salivary gland

## **Blood Group Antigen Precursor Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Blood Group Antigen Precursor Polyclonal Antibody - Images**