

**G protein beta subunit like Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58901**

## Specification

---

### G protein beta subunit like Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	<a href="#">O9BVC4</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35876

### G protein beta subunit like Polyclonal Antibody - Additional Information

Gene ID 64223

#### Other Names

Target of rapamycin complex subunit LST8, TORC subunit LST8, G protein beta subunit-like, Gable, Protein GbetaL, Mammalian lethal with SEC13 protein 8, mLST8, MLST8, GBL, LST8

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### G protein beta subunit like Polyclonal Antibody - Protein Information

**Name** MLST8 {ECO:0000303|PubMed:34741373, ECO:0000312|HGNC:HGNC:24825}

#### Function

Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals (PubMed: <a href="http://www.uniprot.org/citations/12718876" target="\_blank">12718876</a>, PubMed: <a href="http://www.uniprot.org/citations/15268862" target="\_blank">15268862</a>, PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>, PubMed: <a href="http://www.uniprot.org/citations/24403073" target="\_blank">24403073</a>, PubMed: <a href="http://www.uniprot.org/citations/28489822" target="\_blank">28489822</a>). mTORC1 is activated in response to growth factors or amino acids (PubMed: <a href="http://www.uniprot.org/citations/12718876" target="\_blank">12718876</a>, PubMed: <a href="http://www.uniprot.org/citations/15268862" target="\_blank">15268862</a>, PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>, PubMed: <a href="http://www.uniprot.org/citations/24403073" target="\_blank">24403073</a>). In response to nutrients, mTORC1 is recruited to the lysosome membrane and promotes protein, lipid and nucleotide synthesis by phosphorylating several substrates, such as ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) (PubMed: <a href="http://www.uniprot.org/citations/12718876" target="\_blank">12718876</a>).

PubMed: <a href="http://www.uniprot.org/citations/15268862" target="\_blank">15268862</a>, PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>, PubMed: <a href="http://www.uniprot.org/citations/24403073" target="\_blank">24403073</a>). In the same time, it inhibits catabolic pathways by phosphorylating the autophagy initiation components ULK1 and ATG13, as well as transcription factor TFEB, a master regulators of lysosomal biogenesis and autophagy (PubMed: <a href="http://www.uniprot.org/citations/24403073" target="\_blank">24403073</a>). The mTORC1 complex is inhibited in response to starvation and amino acid depletion (PubMed: <a href="http://www.uniprot.org/citations/24403073" target="\_blank">24403073</a>). Within mTORC1, MLST8 interacts directly with MTOR and enhances its kinase activity (PubMed: <a href="http://www.uniprot.org/citations/12718876" target="\_blank">12718876</a>). In nutrient-poor conditions, stabilizes the MTOR- RPTOR interaction and favors RPTOR-mediated inhibition of MTOR activity (PubMed: <a href="http://www.uniprot.org/citations/12718876" target="\_blank">12718876</a>). As part of the mTORC2 complex, transduces signals from growth factors to pathways involved in proliferation, cytoskeletal organization, lipogenesis and anabolic output (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>, PubMed: <a href="http://www.uniprot.org/citations/35926713" target="\_blank">35926713</a>). mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>, PubMed: <a href="http://www.uniprot.org/citations/35926713" target="\_blank">35926713</a>). In response to growth factors, mTORC2 phosphorylates and activates AGC protein kinase family members, including AKT (AKT1, AKT2 and AKT3), PKC (PRKCA, PRKCB and PRKCE) and SGK1 (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>, PubMed: <a href="http://www.uniprot.org/citations/35926713" target="\_blank">35926713</a>). mTORC2 functions upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>). mTORC2 promotes the serum-induced formation of stress-fibers or F-actin (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>). mTORC2 plays a critical role in AKT1 activation by mediating phosphorylation of different sites depending on the context, such as 'Thr-450', 'Ser-473', 'Ser-477' or 'Thr-479', facilitating the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1/PDK1 which is a prerequisite for full activation (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>). mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422' (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>). mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657' (PubMed: <a href="http://www.uniprot.org/citations/15467718" target="\_blank">15467718</a>). Within mTORC2, MLST8 acts as a bridge between MAPKAP1/SIN1 and MTOR (PubMed: <a href="http://www.uniprot.org/citations/31085701" target="\_blank">31085701</a>).

### Cellular Location

Lysosome membrane. Cytoplasm {ECO:0000250|UniProtKB:Q9Z2K5}. Note=Targeting to lysosomal membrane depends on amino acid availability: mTORC1 is recruited to lysosome membranes via interaction with GTP-bound form of RagA/RRAGA (or RagB/RRAGB) in complex with the GDP-bound form of RagC/RRAGC (or RagD/RRAGD), promoting its mTORC1 recruitment to the lysosomes

### Tissue Location

Broadly expressed, with highest levels in skeletal muscle, heart and kidney.


## G protein beta subunit like Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **G protein beta subunit like Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GBL protein) Polyclonal Antibody, Unconjugated (bs-8161R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.