

WDR77 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59053

Specification

WDR77 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O9BQA1
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36724

WDR77 Polyclonal Antibody - Additional Information

Gene ID 79084

Other Names

Methylosome protein 50, MEP-50, Androgen receptor cofactor p44, WD repeat-containing protein 77, p44/Mep50, WDR77, MEP50, WD45

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

WDR77 Polyclonal Antibody - Protein Information

Name WDR77 ([HGNC:29652](#))

Function

Non-catalytic component of the methylosome complex, composed of PRMT5, WDR77 and CLNS1A, which modifies specific arginines to dimethylarginines in several spliceosomal Sm proteins and histones (PubMed:11756452). This modification targets Sm proteins to the survival of motor neurons (SMN) complex for assembly into small nuclear ribonucleoprotein core particles. Might play a role in transcription regulation. The methylosome complex also methylates the Piwi proteins (PIWIL1, PIWIL2 and PIWIL4), methylation of Piwi proteins being required for the interaction with Tudor domain-containing proteins and subsequent localization to the meiotic nuage (PubMed:23071334).

Cellular Location

Nucleus. Cytoplasm. Note=Nuclear in Leydig cells and cytoplasmic in germ cells during fetal testicular development. In adult testis, predominantly nuclear. Subcellular location varies from nuclear to cytoplasmic in various tumors (PubMed:17437848).

Tissue Location

Highly expressed in heart, skeletal muscle, spleen, testis, uterus, prostate and thymus. In testis, expressed in germ cells and Leydig cells, but not in peritubular myocytes, nor in Sertoli cells. Expressed in prostate cancers, in seminomas and in Leydig cell tumors.

WDR77 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDR77 Polyclonal Antibody - Images