

NUDT21 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59083

Specification

NUDT21 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O43809
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26227

NUDT21 Polyclonal Antibody - Additional Information

Gene ID 11051

Other Names

Cleavage and polyadenylation specificity factor subunit 5, Cleavage and polyadenylation specificity factor 25 kDa subunit, CPSF 25 kDa subunit, Cleavage factor Im complex 25 kDa subunit, CFIm25, Nucleoside diphosphate-linked moiety X motif 21, Nudix motif 21, Nudix hydrolase 21, Pre-mRNA cleavage factor Im 68 kDa subunit, NUDT21 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=13870)
target="_blank">HGNC:13870

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NUDT21 Polyclonal Antibody - Protein Information

Name NUDT21 ([HGNC:13870](#))

Function

Component of the cleavage factor Im (CFIm) complex that functions as an activator of the pre-mRNA 3'-end cleavage and polyadenylation processing required for the maturation of pre-mRNA into functional mRNAs (PubMed: [14690600](http://www.uniprot.org/citations/14690600), PubMed: [15937220](http://www.uniprot.org/citations/15937220), PubMed: [17024186](http://www.uniprot.org/citations/17024186), PubMed: [17098938](http://www.uniprot.org/citations/17098938), PubMed: [29276085](http://www.uniprot.org/citations/29276085), PubMed: [8626397](http://www.uniprot.org/citations/8626397), PubMed: [9659921](http://www.uniprot.org/citations/9659921)). CFIm contributes to the recruitment of multiprotein complexes on specific sequences on the pre-mRNA 3'-end, so called cleavage and polyadenylation signals (pA

signals) (PubMed:14690600, PubMed:17024186, PubMed:8626397, PubMed:9659921). Most pre-mRNAs contain multiple pA signals, resulting in alternative cleavage and polyadenylation (APA) producing mRNAs with variable 3'-end formation (PubMed:17098938, PubMed:23187700, PubMed:29276085). The CFIm complex acts as a key regulator of cleavage and polyadenylation site choice during APA through its binding to 5'-UGUA-3' elements localized in the 3'- untranslated region (UTR) for a huge number of pre-mRNAs (PubMed:17098938, PubMed:20695905, PubMed:29276085). NUDT21/CPSF5 activates indirectly the mRNA 3'-processing machinery by recruiting CPSF6 and/or CPSF7 (PubMed:29276085). Binds to 5'-UGUA-3' elements localized upstream of pA signals that act as enhancers of pre-mRNA 3'- end processing (PubMed:14690600, PubMed:15169763, PubMed:17024186, PubMed:20479262, PubMed:22813749, PubMed:8626397). The homodimer mediates simultaneous sequence-specific recognition of two 5'-UGUA-3' elements within the pre-mRNA (PubMed:20479262, PubMed:21295486). Plays a role in somatic cell fate transitions and pluripotency by regulating widespread changes in gene expression through an APA-dependent function (By similarity). Binds to chromatin (By similarity). Binds to, but does not hydrolyze mono- and di-adenosine nucleotides (PubMed:18445629).

Cellular Location

Nucleus. Cytoplasm Note=Shuttles between the nucleus and the cytoplasm in a transcription- and XPO1/CRM1-independent manner, most probably in complex with the cleavage factor Im complex (CFIm) (PubMed:19864460) In punctate subnuclear structures localized adjacent to nuclear speckles, called paraspeckles (PubMed:15169763)

Tissue Location

Expressed in the heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

NUDT21 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NUDT21 Polyclonal Antibody - Images