

**OIP-5/CT86 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP59088**

## Specification

---

### OIP-5/CT86 Polyclonal Antibody - Product Information

Application	<b>WB, IHC-P, IHC-F, IF, ICC, E</b>
Primary Accession	<a href="#">O43482</a>
Reactivity	<b>Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>24691</b>

### OIP-5/CT86 Polyclonal Antibody - Additional Information

**Gene ID** 11339

#### Other Names

Protein Mis18-beta, Cancer/testis antigen 86, CT86, Opa-interacting protein 5, OIP-5, OIP5 {ECO:0000303|PubMed:9466265}, MIS18B

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### OIP-5/CT86 Polyclonal Antibody - Protein Information

**Name** OIP5 {ECO:0000303|PubMed:9466265}

**Synonyms** MIS18B

#### Function

Required for recruitment of CENPA to centromeres and normal chromosome segregation during mitosis.

#### Cellular Location

Nucleus. Chromosome. Chromosome, centromere. Note=Associated with centromeres in interphase cells, from late anaphase to the G1 phase. Not detected on centromeres during earlier phases of mitosis. Associated with chromatin

### OIP-5/CT86 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**OIP-5/CT86 Polyclonal Antibody - Images**