

RNF167 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59189

Specification

RNF167 Polyclonal Antibody - Product Information

Application	IHC-F, IF, E
Primary Accession	O9H6Y7
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38299

RNF167 Polyclonal Antibody - Additional Information

Gene ID 26001

Other Names

E3 ubiquitin-protein ligase RNF167, 2.3.2.27, RING finger protein 167, RING-type E3 ubiquitin transferase RNF167, RING105, RNF167

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RNF167 Polyclonal Antibody - Protein Information

Name RNF167 {ECO:0000303|PubMed:23353890, ECO:0000312|HGNC:HGNC:24544}

Function

E3 ubiquitin-protein ligase that acts as a regulator of the TORC1 signaling pathway (PubMed:[33594058](http://www.uniprot.org/citations/33594058), PubMed:[35114100](http://www.uniprot.org/citations/35114100)). Positively regulates the TORC1 signaling pathway independently of arginine levels: acts by catalyzing 'Lys-29'-polyubiquitination and degradation of CASTOR1, releasing the GATOR2 complex from CASTOR1 (PubMed:[33594058](http://www.uniprot.org/citations/33594058)). Also negatively regulates the TORC1 signaling pathway in response to leucine deprivation: acts by mediating 'Lys-63'-linked polyubiquitination of SESN2, promoting SESN2-interaction with the GATOR2 complex (PubMed:[35114100](http://www.uniprot.org/citations/35114100)). Also involved in protein trafficking and localization (PubMed:[23129617](http://www.uniprot.org/citations/23129617), PubMed:[23353890](http://www.uniprot.org/citations/23353890), PubMed:[24387786](http://www.uniprot.org/citations/24387786), PubMed:[27808481](http://www.uniprot.org/citations/27808481), PubMed:[27808481](http://www.uniprot.org/citations/27808481)).

<http://www.uniprot.org/citations/32409562> target="_blank">32409562). Acts as a regulator of synaptic transmission by mediating ubiquitination and degradation of AMPAR receptor GluA2/GRIA2 (PubMed:23129617, PubMed:33650289). Does not catalyze ubiquitination of GluA1/GRIA1 (PubMed:23129617). Also acts as a regulator of the recycling endosome pathway by mediating ubiquitination of VAMP3 (PubMed:23353890). Regulates lysosome positioning by catalyzing ubiquitination and degradation of ARL8B (PubMed:27808481). Plays a role in growth regulation involved in G1/S transition by mediating, possibly by mediating ubiquitination of SLC22A18 (PubMed:16314844). Acts with a limited set of E2 enzymes, such as UBE2D1 and UBE2N (PubMed:33650289).

Cellular Location

Lysosome membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Endomembrane system; Single-pass membrane protein. Cell membrane; Single-pass type I membrane protein. Note=Targeted to cytoplasmic membranes; mainly localizes to lysosomal membrane (PubMed:16314844, PubMed:23129617). A subpopulation localizes to the cell membrane of neurons (PubMed:23129617). [Isoform 2]: Cytoplasm, cytosol

Tissue Location

Widely expressed (PubMed:23129617). Strongly expressed in the kidney, pancreas, testis and liver (at protein level) (PubMed:16314844, PubMed:23129617).

RNF167 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RNF167 Polyclonal Antibody - Images