

C17orf75 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59324

Specification

C17orf75 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O9HAS0
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44622

C17orf75 Polyclonal Antibody - Additional Information

Gene ID 64149

Other Names

Protein Njmu-R1, C17orf75

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C17orf75 Polyclonal Antibody - Protein Information

Name C17orf75

Function

As component of the WDR11 complex acts together with TBC1D23 to facilitate the golgin-mediated capture of vesicles generated using AP-1 (PubMed:29426865). May have a role in spermatogenesis.

Cellular Location

Golgi apparatus, trans-Golgi network. Cytoplasmic vesicle

Tissue Location


Highly expressed in testis and also expressed in fetal testis

C17orf75 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C17orf75 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C17orf75) Polyclonal Antibody, Unconjugated (bs-9646R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Sample:

U2os Cell (Human) Lysate at 30 ug
Primary: Anti- C17orf75 (bs-9646R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 45kD
Observed band size: 45kD