

Anti-CDC34 Antibody
Rabbit polyclonal antibody to CDC34
Catalog # AP59507

Specification

Anti-CDC34 Antibody - Product Information

Application	WB
Primary Accession	P49427
Other Accession	Q8CFI2
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26737

Anti-CDC34 Antibody - Additional Information

Gene ID 997

Other Names

UBCH3; UBE2R1; Ubiquitin-conjugating enzyme E2 R1; Ubiquitin-conjugating enzyme E2-32 kDa complementing; Ubiquitin-conjugating enzyme E2-CDC34; Ubiquitin-protein ligase R1

Target/Specificity

Recognizes endogenous levels of CDC34 protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-CDC34 Antibody - Protein Information

Name CDC34

Synonyms UBCH3, UBE2R1

Function

Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-48'- linked polyubiquitination (PubMed:22496338). Cooperates with the E2 UBCH5C and the SCF(FBXW11) E3 ligase complex for the polyubiquitination of NFKBIA leading to its subsequent proteasomal degradation. Performs ubiquitin chain elongation building ubiquitin chains from the UBE2D3- primed NFKBIA-linked ubiquitin. UBE2D3 acts as an initiator E2,

priming the phosphorylated NFKBIA target at positions 'Lys-21' and/or 'Lys-22' with a monoubiquitin. Cooperates with the SCF(SKP2) E3 ligase complex to regulate cell proliferation through ubiquitination and degradation of MYBL2 and KIP1. Involved in ubiquitin conjugation and degradation of CREM isoform ICERIIgamma and ATF15 resulting in abrogation of ICERIIgamma and ATF5-mediated repression of cAMP-induced transcription during both meiotic and mitotic cell cycles. Involved in the regulation of the cell cycle G2/M phase through its targeting of the WEE1 kinase for ubiquitination and degradation. Also involved in the degradation of beta-catenin. Is target of human herpes virus 1 protein ICP0, leading to ICP0-dependent dynamic interaction with proteasomes (PubMed: 10329681, PubMed: 10373550, PubMed: 10871850, PubMed: 11675391, PubMed: 12037680, PubMed: 15652359, PubMed: 17461777, PubMed: 17698585, PubMed: 19112177, PubMed: 19126550, PubMed: 19945379, PubMed: 20061386, PubMed: 20347421).

Cellular Location

Cytoplasm. Nucleus. Note=The phosphorylation of the C-terminal tail plays an important role in mediating nuclear localization. Colocalizes with beta-tubulin on mitotic spindles in anaphase

Tissue Location

Expressed in testes during spermatogenesis to regulate repression of cAMP-induced transcription


Anti-CDC34 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CDC34 Antibody - Images





Western blot analysis of CDC34 expression in mouse lung (A), mouse liver (B) whole cell lysates.

Anti-CDC34 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDC34. The exact sequence is proprietary.