

**Anti-HADH2 Antibody**  
Rabbit polyclonal antibody to HADH2  
Catalog # AP59577

## Specification

---

### Anti-HADH2 Antibody - Product Information

Application	WB, IF
Primary Accession	<a href="#">O99714</a>
Other Accession	<a href="#">O08756</a>
Reactivity	Human, Mouse, Rat, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26923

### Anti-HADH2 Antibody - Additional Information

Gene ID 3028

#### Other Names

ERAB; HADH2; MRPP2; SCHAD; XH98G2; 3-hydroxyacyl-CoA dehydrogenase type-2; 17-beta-hydroxysteroid dehydrogenase 10; 17-beta-HSD 10; 3-hydroxy-2-methylbutyryl-CoA dehydrogenase; 3-hydroxyacyl-CoA dehydrogenase type II; Endoplasmic reticulum-associated amyloid beta-peptide-binding protein; Mitochondrial ribonuclease P protein 2; Mitochondrial RNase P protein 2; Short-chain type dehydrogenase/reductase XH98G2; Type II HADH

#### Target/Specificity

Recognizes endogenous levels of HADH2 protein.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)  
IF~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### Anti-HADH2 Antibody - Protein Information

Name HSD17B10

Synonyms ERAB, HADH2, MRPP2, SCHAD, SDR5C1, XH98G

#### Function

Mitochondrial dehydrogenase involved in pathways of fatty acid, branched-chain amino acid and steroid metabolism (PubMed: <http://www.uniprot.org/citations/10600649>)

target="\_blank">10600649</a>, PubMed:<a href="http://www.uniprot.org/citations/12917011" target="\_blank">12917011</a>, PubMed:<a href="http://www.uniprot.org/citations/18996107" target="\_blank">18996107</a>, PubMed:<a href="http://www.uniprot.org/citations/19706438" target="\_blank">19706438</a>, PubMed:<a href="http://www.uniprot.org/citations/20077426" target="\_blank">20077426</a>, PubMed:<a href="http://www.uniprot.org/citations/25925575" target="\_blank">25925575</a>, PubMed:<a href="http://www.uniprot.org/citations/26950678" target="\_blank">26950678</a>, PubMed:<a href="http://www.uniprot.org/citations/28888424" target="\_blank">28888424</a>, PubMed:<a href="http://www.uniprot.org/citations/9553139" target="\_blank">9553139</a>). Acts as (S)-3-hydroxyacyl-CoA dehydrogenase in mitochondrial fatty acid beta-oxidation, a major degradation pathway of fatty acids. Catalyzes the third step in the beta-oxidation cycle, namely the reversible conversion of (S)-3-hydroxyacyl-CoA to 3-ketoacyl-CoA. Preferentially accepts straight medium- and short-chain acyl-CoA substrates with highest efficiency for (3S)-hydroxybutanoyl-CoA (PubMed:<a href="http://www.uniprot.org/citations/10600649" target="\_blank">10600649</a>, PubMed:<a href="http://www.uniprot.org/citations/12917011" target="\_blank">12917011</a>, PubMed:<a href="http://www.uniprot.org/citations/25925575" target="\_blank">25925575</a>, PubMed:<a href="http://www.uniprot.org/citations/26950678" target="\_blank">26950678</a>, PubMed:<a href="http://www.uniprot.org/citations/9553139" target="\_blank">9553139</a>). Acts as 3-hydroxy-2-methylbutyryl-CoA dehydrogenase in branched-chain amino acid catabolic pathway. Catalyzes the oxidation of 3-hydroxy-2-methylbutanoyl-CoA into 2-methyl-3-oxobutanoyl-CoA, a step in isoleucine degradation pathway (PubMed:<a href="http://www.uniprot.org/citations/18996107" target="\_blank">18996107</a>, PubMed:<a href="http://www.uniprot.org/citations/19706438" target="\_blank">19706438</a>, PubMed:<a href="http://www.uniprot.org/citations/20077426" target="\_blank">20077426</a>). Has hydroxysteroid dehydrogenase activity toward steroid hormones and bile acids. Catalyzes the oxidation of 3alpha-, 17beta-, 20beta- and 21- hydroxysteroids and 7alpha- and 7beta-hydroxy bile acids (PubMed:<a href="http://www.uniprot.org/citations/10600649" target="\_blank">10600649</a>, PubMed:<a href="http://www.uniprot.org/citations/12917011" target="\_blank">12917011</a>). Oxidizes allopregnanolone/brexanolone at the 3alpha-hydroxyl group, which is known to be critical for the activation of gamma-aminobutyric acid receptors (GABAARs) chloride channel (PubMed:<a href="http://www.uniprot.org/citations/19706438" target="\_blank">19706438</a>, PubMed:<a href="http://www.uniprot.org/citations/28888424" target="\_blank">28888424</a>). Has phospholipase C-like activity toward cardiolipin and its oxidized species. Likely oxidizes the 2'-hydroxyl in the head group of cardiolipin to form a ketone intermediate that undergoes nucleophilic attack by water and fragments into diacylglycerol, dihydroxyacetone and orthophosphate. Has higher affinity for cardiolipin with oxidized fatty acids and may degrade these species during the oxidative stress response to protect cells from apoptosis (PubMed:<a href="http://www.uniprot.org/citations/26338420" target="\_blank">26338420</a>). By interacting with intracellular amyloid-beta, it may contribute to the neuronal dysfunction associated with Alzheimer disease (AD) (PubMed:<a href="http://www.uniprot.org/citations/9338779" target="\_blank">9338779</a>). Essential for structural and functional integrity of mitochondria (PubMed:<a href="http://www.uniprot.org/citations/20077426" target="\_blank">20077426</a>).

### Cellular Location

Mitochondrion. Mitochondrion matrix, mitochondrion nucleoid

### Tissue Location

Ubiquitously expressed in normal tissues but is overexpressed in neurons affected in AD.


### Anti-HADH2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-HADH2 Antibody - Images



Western blot analysis of HADH2 expression in H446 (A), U2OS (B), DLD (C), mouse lung (D), mouse liver (E) whole cell lysates.



Immunohistochemical analysis of HADH2 staining in human lung formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of HADH2 staining in SKNSH cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-HADH2 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HADH2. The exact sequence is proprietary.