

**Anti-CD51 HC Antibody**  
Rabbit polyclonal antibody to CD51 HC  
Catalog # AP59595

## Specification

---

### Anti-CD51 HC Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P06756</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116038

### Anti-CD51 HC Antibody - Additional Information

Gene ID 3685

#### Other Names

MSK8; VNRA; Integrin alpha-V; Vitronectin receptor subunit alpha; CD51

#### Target/Specificity

Recognizes endogenous levels of CD51 HC protein.

#### Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### Anti-CD51 HC Antibody - Protein Information

Name ITGAV ([HGNC:6150](#))

#### Function

The alpha-V (ITGAV) integrins are receptors for vitronectin, cytotactin, fibronectin, fibrinogen, laminin, matrix metalloproteinase- 2, osteopontin, osteomodulin, prothrombin, thrombospondin and vWF. They recognize the sequence R-G-D in a wide array of ligands. ITGAV:ITGB3 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1- dependent fractalkine signaling (PubMed:<a href="http://www.uniprot.org/citations/23125415" target="\_blank">23125415</a>). ITGAV:ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:<a href="http://www.uniprot.org/citations/20682778" target="\_blank">20682778</a>). ITGAV:ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling (PubMed:<a href="http://www.uniprot.org/citations/18441324" target="\_blank">18441324</a>). ITGAV:ITGB3 binds to FGF2 and this binding is essential for

FGF2 signaling (PubMed:<a href="http://www.uniprot.org/citations/28302677" target="\_blank">28302677</a>). ITGAV:ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:<a href="http://www.uniprot.org/citations/19578119" target="\_blank">19578119</a>). ITGAV:ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:<a href="http://www.uniprot.org/citations/28873464" target="\_blank">28873464</a>). ITGAV:ITGB3 binds to IL1B and this binding is essential for IL1B signaling (PubMed:<a href="http://www.uniprot.org/citations/29030430" target="\_blank">29030430</a>). ITGAV:ITGB3 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:<a href="http://www.uniprot.org/citations/18635536" target="\_blank">18635536</a>, PubMed:<a href="http://www.uniprot.org/citations/25398877" target="\_blank">25398877</a>). ITGAV:ITGB3 and ITGAV:ITGB6 act as receptors for fibrillin-1 (FBN1) and mediate R-G-D-dependent cell adhesion to FBN1 (PubMed:<a href="http://www.uniprot.org/citations/12807887" target="\_blank">12807887</a>, PubMed:<a href="http://www.uniprot.org/citations/17158881" target="\_blank">17158881</a>). Integrin alpha-V/beta-6 or alpha-V/beta-8 (ITGAV:ITGB6 or ITGAV:ITGB8) mediates R-G-D-dependent release of transforming growth factor beta-1 (TGF-beta-1) from regulatory Latency-associated peptide (LAP), thereby playing a key role in TGF-beta-1 activation (PubMed:<a href="http://www.uniprot.org/citations/15184403" target="\_blank">15184403</a>, PubMed:<a href="http://www.uniprot.org/citations/22278742" target="\_blank">22278742</a>, PubMed:<a href="http://www.uniprot.org/citations/28117447" target="\_blank">28117447</a>). ITGAV:ITGB3 acts as a receptor for CD40LG (PubMed:<a href="http://www.uniprot.org/citations/31331973" target="\_blank">31331973</a>). ITGAV:ITGB3 acts as a receptor for IBSP and promotes cell adhesion and migration to IBSP (PubMed:<a href="http://www.uniprot.org/citations/10640428" target="\_blank">10640428</a>).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell junction, focal adhesion


#### Anti-CD51 HC Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-CD51 HC Antibody - Images





Western blot analysis of CD51 HC expression in A549 (A), HeLa (B), C6 (C), MEF (D) whole cell lysates.



Immunohistochemical analysis of CD51 HC staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-CD51 HC Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD51 HC. The exact sequence is proprietary.