

Anti-LIMK1/2 (pT508/505) Antibody
Rabbit polyclonal antibody to LIMK1/2 (pT508/505)
Catalog # AP59607

Specification

Anti-LIMK1/2 (pT508/505) Antibody - Product Information

Application	WB, IF
Primary Accession	P53667
Reactivity	Human, Rat, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	72585

Anti-LIMK1/2 (pT508/505) Antibody - Additional Information

Gene ID 3984

Other Names

LIMK; LIM domain kinase 1; LIMK-1

Target/Specificity

Recognizes endogenous levels of LIMK1/2 (pT508/505) protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)
IF~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-LIMK1/2 (pT508/505) Antibody - Protein Information

Name LIMK1

Synonyms LIMK

Function

Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways
(PubMed: [10436159](http://www.uniprot.org/citations/10436159)),
PubMed: [11832213](http://www.uniprot.org/citations/11832213),
PubMed: [12807904](http://www.uniprot.org/citations/12807904),
PubMed: [15660133](http://www.uniprot.org/citations/15660133),
PubMed: [16230460](http://www.uniprot.org/citations/16230460),

PubMed: [18028908](http://www.uniprot.org/citations/18028908), PubMed: [22328514](http://www.uniprot.org/citations/22328514), PubMed: [23633677](http://www.uniprot.org/citations/23633677)). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed: [10436159](http://www.uniprot.org/citations/10436159)). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destrin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed: [11832213](http://www.uniprot.org/citations/11832213), PubMed: [15660133](http://www.uniprot.org/citations/15660133), PubMed: [16230460](http://www.uniprot.org/citations/16230460), PubMed: [23633677](http://www.uniprot.org/citations/23633677)). In this way LIMK1 regulates several actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed: [11832213](http://www.uniprot.org/citations/11832213), PubMed: [15660133](http://www.uniprot.org/citations/15660133), PubMed: [16230460](http://www.uniprot.org/citations/16230460), PubMed: [23633677](http://www.uniprot.org/citations/23633677)). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed: [18028908](http://www.uniprot.org/citations/18028908)). Stimulates axonal outgrowth and may be involved in brain development (PubMed: [18028908](http://www.uniprot.org/citations/18028908)).

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P53668} Note=Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner. {ECO:0000250|UniProtKB:P53668}

Tissue Location

Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle


Anti-LIMK1/2 (pT508/505) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-LIMK1/2 (pT508/505) Antibody - Images






Western blot analysis of LIMK1/2 (pT508/505) expression in rat brain (A), HEK293T-PMA-15min (B), HEK293T-PMA-5min (C), HEK293T (D) whole cell lysates.



Immunohistochemical analysis of LIMK1/2 (pT508/505) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of LIMK1/2 (pT508/505) staining in PC12 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in

the dark.

Anti-LIMK1/2 (pT508/505) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human LIMK1/2. The exact sequence is proprietary.