

Anti-NSF Antibody
Rabbit polyclonal antibody to NSF
Catalog # AP59643

Specification

Anti-NSF Antibody - Product Information

Application	WB
Primary Accession	P46459
Other Accession	P46460
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	82594

Anti-NSF Antibody - Additional Information

Gene ID 4905

Other Names

Vesicle-fusing ATPase; N-ethylmaleimide-sensitive fusion protein; NEM-sensitive fusion protein; Vesicular-fusion protein NSF

Target/Specificity

Recognizes endogenous levels of NSF protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-NSF Antibody - Protein Information

Name NSF

Function

Required for vesicle-mediated transport. Catalyzes the fusion of transport vesicles within the Golgi cisternae. Is also required for transport from the endoplasmic reticulum to the Golgi stack. Seems to function as a fusion protein required for the delivery of cargo proteins to all compartments of the Golgi stack independent of vesicle origin. Interaction with AMPAR subunit GRIA2 leads to influence GRIA2 membrane cycling (By similarity).

Cellular Location


Cytoplasm.

Anti-NSF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-NSF Antibody - Images



Western blot analysis of NSF expression in HGC27 (A) whole cell lysates.

Anti-NSF Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NSF. The exact sequence is proprietary.