

Anti-Pepsinogen C Antibody
Rabbit polyclonal antibody to Pepsinogen C
Catalog # AP59659

Specification

Anti-Pepsinogen C Antibody - Product Information

Application	WB
Primary Accession	P20142
Other Accession	Q9D7R7
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42426

Anti-Pepsinogen C Antibody - Additional Information

Gene ID 5225

Other Names
Gastricsin; Pepsinogen C

Target/Specificity
Recognizes endogenous levels of Pepsinogen C protein.

Dilution
WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

Format
Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage
Store at -20 °C. Stable for 12 months from date of receipt

Anti-Pepsinogen C Antibody - Protein Information

Name PGC

Function
Hydrolyzes a variety of proteins.


Cellular Location
Secreted.

Anti-Pepsinogen C Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Pepsinogen C Antibody - Images



Western blot analysis of Pepsinogen C expression in HepG2 (A), LO2 (B), SGC7901 (C), AML12 (D), PC12 (E) whole cell lysates.



Immunohistochemical analysis of Pepsinogen C staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. AEC was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-Pepsinogen C Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Pepsinogen C. The exact sequence is proprietary.