

**Anti-CDC2 (pY15) Antibody**  
Rabbit polyclonal antibody to CDC2 (pY15)  
Catalog # AP59982

## Specification

---

### Anti-CDC2 (pY15) Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P06493</a>
Other Accession	<a href="#">P11440</a>
Reactivity	Human, Mouse, Rat, Zebrafish, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34095

### Anti-CDC2 (pY15) Antibody - Additional Information

Gene ID 983

#### Other Names

CDC2; CDC28A; CDKN1; P34CDC2; Cyclin-dependent kinase 1; CDK1; Cell division control protein 2 homolog; Cell division protein kinase 1; p34 protein kinase

#### Target/Specificity

Recognizes endogenous levels of CDC2 (pY15) protein.

#### Dilution

WB~~WB (1/500 - 1/1000)

#### Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### Anti-CDC2 (pY15) Antibody - Protein Information

Name CDK1

Synonyms CDC2, CDC28A, CDKN1, P34CDC2

#### Function

Plays a key role in the control of the eukaryotic cell cycle by modulating the centrosome cycle as well as mitotic onset; promotes G2-M transition via association with multiple interphase cyclins (PubMed: [16407259](http://www.uniprot.org/citations/16407259), PubMed: [16933150](http://www.uniprot.org/citations/16933150), PubMed: [17459720](http://www.uniprot.org/citations/17459720)),

PubMed: <a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>,  
PubMed: <a href="http://www.uniprot.org/citations/19509060" target="\_blank">19509060</a>,  
PubMed: <a href="http://www.uniprot.org/citations/19917720" target="\_blank">19917720</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20171170" target="\_blank">20171170</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20935635" target="\_blank">20935635</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20937773" target="\_blank">20937773</a>,  
PubMed: <a href="http://www.uniprot.org/citations/21063390" target="\_blank">21063390</a>,  
PubMed: <a href="http://www.uniprot.org/citations/2188730" target="\_blank">2188730</a>,  
PubMed: <a href="http://www.uniprot.org/citations/23355470" target="\_blank">23355470</a>,  
PubMed: <a href="http://www.uniprot.org/citations/2344612" target="\_blank">2344612</a>,  
PubMed: <a href="http://www.uniprot.org/citations/23601106" target="\_blank">23601106</a>,  
PubMed: <a href="http://www.uniprot.org/citations/23602554" target="\_blank">23602554</a>,  
PubMed: <a href="http://www.uniprot.org/citations/25556658" target="\_blank">25556658</a>,  
PubMed: <a href="http://www.uniprot.org/citations/26829474" target="\_blank">26829474</a>,  
PubMed: <a href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491</a>,  
PubMed: <a href="http://www.uniprot.org/citations/30139873" target="\_blank">30139873</a>,  
PubMed: <a href="http://www.uniprot.org/citations/30704899" target="\_blank">30704899</a>).  
Phosphorylates PARVA/actopaxin, APC, AMPH, APC, BARD1, Bcl-xL/BCL2L1, BRCA2, CALD1, CASP8,  
CDC7, CDC20, CDC25A, CDC25C, CC2D1A, CENPA, CSNK2 proteins/CKII, FZR1/CDH1, CDK7,  
CEBPB, CHAMP1, DMD/dystrophin, EEF1 proteins/EF-1, EZH2, KIF11/EG5, EGFR, FANCG, FOS, GFAP,  
GOLGA2/GM130, GRASP1, UBE2A/hHR6A, HIST1H1 proteins/histone H1, HMGA1, HIVEP3/KRC,  
KAT5, LMNA, LMNB, LBR, MKI67, LATS1, MAP1B, MAP4, MARCKS, MCM2, MCM4, MKLP1, MLST8,  
MYB, NEFH, NFIC, NPC/nuclear pore complex, PITPNM1/NIR2, NPM1, NCL, NUCKS1, NPM1/numatrin,  
ORC1, PRKAR2A, EEF1E1/p18, EIF3F/p47, p53/TP53, NONO/p54NRB, PAPOLA, PLEC/plectin, RB1,  
TPPP, UL40/R2, RAB4A, RAP1GAP, RBBP8/CtIP, RCC1, RPS6KB1/S6K1, KHDRBS1/SAM68, ESPL1, SKI,  
BIRC5/survivin, STIP1, TEX14, beta-tubulins, MAPT/TAU, NEDD1, VIM/vimentin, TK1, FOXO1,  
RUNX1/AML1, SAMHD1, SIRT2, CGAS and RUNX2 (PubMed: <a href="http://www.uniprot.org/citations/16407259" target="\_blank">16407259</a>,  
PubMed: <a href="http://www.uniprot.org/citations/16933150" target="\_blank">16933150</a>,  
PubMed: <a href="http://www.uniprot.org/citations/17459720" target="\_blank">17459720</a>,  
PubMed: <a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>,  
PubMed: <a href="http://www.uniprot.org/citations/19202191" target="\_blank">19202191</a>,  
PubMed: <a href="http://www.uniprot.org/citations/19509060" target="\_blank">19509060</a>,  
PubMed: <a href="http://www.uniprot.org/citations/19917720" target="\_blank">19917720</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20171170" target="\_blank">20171170</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20935635" target="\_blank">20935635</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20937773" target="\_blank">20937773</a>,  
PubMed: <a href="http://www.uniprot.org/citations/21063390" target="\_blank">21063390</a>,  
PubMed: <a href="http://www.uniprot.org/citations/2188730" target="\_blank">2188730</a>,  
PubMed: <a href="http://www.uniprot.org/citations/23355470" target="\_blank">23355470</a>,  
PubMed: <a href="http://www.uniprot.org/citations/2344612" target="\_blank">2344612</a>,  
PubMed: <a href="http://www.uniprot.org/citations/23601106" target="\_blank">23601106</a>,  
PubMed: <a href="http://www.uniprot.org/citations/23602554" target="\_blank">23602554</a>,  
PubMed: <a href="http://www.uniprot.org/citations/25012651" target="\_blank">25012651</a>,  
PubMed: <a href="http://www.uniprot.org/citations/25556658" target="\_blank">25556658</a>,  
PubMed: <a href="http://www.uniprot.org/citations/26829474" target="\_blank">26829474</a>,  
PubMed: <a href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491</a>,  
PubMed: <a href="http://www.uniprot.org/citations/30704899" target="\_blank">30704899</a>,  
PubMed: <a href="http://www.uniprot.org/citations/32351706" target="\_blank">32351706</a>,  
PubMed: <a href="http://www.uniprot.org/citations/34741373" target="\_blank">34741373</a>).  
CDK1/CDC2-cyclin-B controls pronuclear union in interphase fertilized eggs (PubMed: <a href="http://www.uniprot.org/citations/18480403" target="\_blank">18480403</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>). Essential for  
early stages of embryonic development (PubMed: <a href="http://www.uniprot.org/citations/18480403" target="\_blank">18480403</a>,  
PubMed: <a href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>). During G2  
and early mitosis, CDC25A/B/C-mediated dephosphorylation activates CDK1/cyclin complexes

which phosphorylate several substrates that trigger at least centrosome separation, Golgi dynamics, nuclear envelope breakdown and chromosome condensation (PubMed:<a href="http://www.uniprot.org/citations/18480403" target="\_blank">18480403</a>, PubMed:<a href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>, PubMed:<a href="http://www.uniprot.org/citations/2188730" target="\_blank">2188730</a>, PubMed:<a href="http://www.uniprot.org/citations/2344612" target="\_blank">2344612</a>, PubMed:<a href="http://www.uniprot.org/citations/30139873" target="\_blank">30139873</a>). Once chromosomes are condensed and aligned at the metaphase plate, CDK1 activity is switched off by WEE1- and PKMYT1-mediated phosphorylation to allow sister chromatid separation, chromosome decondensation, reformation of the nuclear envelope and cytokinesis (PubMed:<a href="http://www.uniprot.org/citations/18480403" target="\_blank">18480403</a>, PubMed:<a href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>). Phosphorylates KRT5 during prometaphase and metaphase (By similarity). Inactivated by PKR/EIF2AK2- and WEE1-mediated phosphorylation upon DNA damage to stop cell cycle and genome replication at the G2 checkpoint thus facilitating DNA repair (PubMed:<a href="http://www.uniprot.org/citations/20360007" target="\_blank">20360007</a>). Reactivated after successful DNA repair through WIP1-dependent signaling leading to CDC25A/B/C-mediated dephosphorylation and restoring cell cycle progression (PubMed:<a href="http://www.uniprot.org/citations/20395957" target="\_blank">20395957</a>). Catalyzes lamin (LMNA, LMNB1 and LMNB2) phosphorylation at the onset of mitosis, promoting nuclear envelope breakdown (PubMed:<a href="http://www.uniprot.org/citations/2188730" target="\_blank">2188730</a>, PubMed:<a href="http://www.uniprot.org/citations/2344612" target="\_blank">2344612</a>, PubMed:<a href="http://www.uniprot.org/citations/37788673" target="\_blank">37788673</a>). In proliferating cells, CDK1-mediated FOXO1 phosphorylation at the G2-M phase represses FOXO1 interaction with 14-3-3 proteins and thereby promotes FOXO1 nuclear accumulation and transcription factor activity, leading to cell death of postmitotic neurons (PubMed:<a href="http://www.uniprot.org/citations/18356527" target="\_blank">18356527</a>). The phosphorylation of beta-tubulins regulates microtubule dynamics during mitosis (PubMed:<a href="http://www.uniprot.org/citations/16371510" target="\_blank">16371510</a>). NEDD1 phosphorylation promotes PLK1-mediated NEDD1 phosphorylation and subsequent targeting of the gamma-tubulin ring complex (gTuRC) to the centrosome, an important step for spindle formation (PubMed:<a href="http://www.uniprot.org/citations/19509060" target="\_blank">19509060</a>). In addition, CC2D1A phosphorylation regulates CC2D1A spindle pole localization and association with SCC1/RAD21 and centriole cohesion during mitosis (PubMed:<a href="http://www.uniprot.org/citations/20171170" target="\_blank">20171170</a>). The phosphorylation of Bcl-xL/BCL2L1 after prolonged G2 arrest upon DNA damage triggers apoptosis (PubMed:<a href="http://www.uniprot.org/citations/19917720" target="\_blank">19917720</a>). In contrast, CASP8 phosphorylation during mitosis prevents its activation by proteolysis and subsequent apoptosis (PubMed:<a href="http://www.uniprot.org/citations/20937773" target="\_blank">20937773</a>). This phosphorylation occurs in cancer cell lines, as well as in primary breast tissues and lymphocytes (PubMed:<a href="http://www.uniprot.org/citations/20937773" target="\_blank">20937773</a>). EZH2 phosphorylation promotes H3K27me3 maintenance and epigenetic gene silencing (PubMed:<a href="http://www.uniprot.org/citations/20935635" target="\_blank">20935635</a>). CALD1 phosphorylation promotes Schwann cell migration during peripheral nerve regeneration (By similarity). CDK1-cyclin-B complex phosphorylates NCKAP5L and mediates its dissociation from centrosomes during mitosis (PubMed:<a href="http://www.uniprot.org/citations/26549230" target="\_blank">26549230</a>). Regulates the amplitude of the cyclic expression of the core clock gene BMAL1 by phosphorylating its transcriptional repressor NR1D1, and this phosphorylation is necessary for SCF(FBXW7)- mediated ubiquitination and proteasomal degradation of NR1D1 (PubMed:<a href="http://www.uniprot.org/citations/27238018" target="\_blank">27238018</a>). Phosphorylates EML3 at 'Thr-881' which is essential for its interaction with HAUS augmin-like complex and TUBG1 (PubMed:<a href="http://www.uniprot.org/citations/30723163" target="\_blank">30723163</a>). Phosphorylates CGAS during mitosis, leading to its inhibition, thereby preventing CGAS activation by self DNA during mitosis (PubMed:<a href="http://www.uniprot.org/citations/32351706" target="\_blank">32351706</a>). Phosphorylates SKA3 on multiple sites during mitosis which

promotes SKA3 binding to the NDC80 complex and anchoring of the SKA complex to kinetochores, to enable stable attachment of mitotic spindle microtubules to kinetochores (PubMed:<a href="http://www.uniprot.org/citations/28479321" target="\_blank">28479321</a>, PubMed:<a href="http://www.uniprot.org/citations/31804178" target="\_blank">31804178</a>, PubMed:<a href="http://www.uniprot.org/citations/32491969" target="\_blank">32491969</a>).

#### Cellular Location

Nucleus {ECO:0000250|UniProtKB:P11440}. Cytoplasm {ECO:0000250|UniProtKB:P11440}. Mitochondrion. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Note=Cytoplasmic during the interphase Colocalizes with SIRT2 on centrosome during prophase and on spindle fibers during metaphase of the mitotic cell cycle. Reversibly translocated from cytoplasm to nucleus when phosphorylated before G2-M transition when associated with cyclin-B1. Accumulates in mitochondria in G2-arrested cells upon DNA-damage

#### Tissue Location


[Isoform 2]: Found in breast cancer tissues.

### Anti-CDC2 (pY15) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-CDC2 (pY15) Antibody - Images



Western blot analysis of CDC2 (pY15) expression in HEK293T (A), HeLa (B) whole cell lysates.

### Anti-CDC2 (pY15) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human CDC2 (pY15). The exact sequence is proprietary.