

**Anti-FGFR1 Antibody**  
**Rabbit polyclonal antibody to FGFR1**  
**Catalog # AP60004**

**Specification**

---

**Anti-FGFR1 Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | <b>WB</b>   |
| Primary Accession | <a href="#">P11362</a>                                    |
| Other Accession   | <a href="#">P16092</a>                                    |
| Reactivity        | <b>Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken</b> |
| Host              | <b>Rabbit</b>   |
| Clonality         | <b>Polyclonal</b>   |
| Calculated MW     | <b>91868</b>  |

**Anti-FGFR1 Antibody - Additional Information**

**Gene ID** 2260

**Other Names**

BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD331

**Target/Specificity**

Recognizes endogenous levels of FGFR1 protein.

**Dilution**

WB~~WB (1/500 - 1/1000)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-FGFR1 Antibody - Protein Information**

**Name** FGFR1

**Synonyms** BFGFR, CEK, FGFBR, FLG, FLT2, HBGFR

**Function**

Tyrosine-protein kinase that acts as a cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of embryonic development, cell proliferation, differentiation and migration. Required for normal mesoderm patterning and correct axial organization during embryonic development, normal skeletogenesis and normal development of the

gonadotropin-releasing hormone (GnRH) neuronal system. Phosphorylates PLCG1, FRS2, GAB1 and SHB. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Promotes phosphorylation of SHC1, STAT1 and PTPN11/SHP2. In the nucleus, enhances RPS6KA1 and CREB1 activity and contributes to the regulation of transcription. FGFR1 signaling is down-regulated by IL17RD/SEF, and by FGFR1 ubiquitination, internalization and degradation.

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. Note=After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus

#### Tissue Location


Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.

### Anti-FGFR1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-FGFR1 Antibody - Images



Western blot analysis of FGFR1 expression in HEK293T (A), HeLa (B) whole cell lysates.

### Anti-FGFR1 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human

FGFR1. The exact sequence is proprietary.