

Anti-ACOT4 Antibody
Rabbit polyclonal antibody to ACOT4
Catalog # AP60125

Specification

Anti-ACOT4 Antibody - Product Information

Application	WB, IF
Primary Accession	O8N9L9
Other Accession	O8BWN8
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46327

Anti-ACOT4 Antibody - Additional Information

Gene ID 122970

Other Names

PTE2B; PTEIB; Acyl-coenzyme A thioesterase 4; Acyl-CoA thioesterase 4; PTE-2b; Peroxisomal acyl coenzyme A thioester hydrolase Ib; Peroxisomal long-chain acyl-CoA thioesterase Ib; PTE-Ib

Target/Specificity

Recognizes endogenous levels of ACOT4 protein.

Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)
IF~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-ACOT4 Antibody - Protein Information

Name ACOT4

Synonyms PTE2B, PTEIB

Function

Catalyzes the hydrolysis of acyl-CoAs into free fatty acids and coenzyme A (CoASH), regulating their respective intracellular levels (PubMed:16940157). Functions as a peroxisomal succinyl-coenzyme A thioesterase that can also hydrolyze glutaryl-CoA and long chain saturated acyl-CoAs (PubMed:16940157).

Cellular Location

Peroxisome.

Tissue Location


Strongest expression in liver and kidney and weaker expression in placenta, heart, and muscle

Anti-ACOT4 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ACOT4 Antibody - Images




Western blot analysis of ACOT4 expression in U2OS (A), MCF7 (B), mouse lung (C), mouse liver (D), rat lung (E), rat liver (F) whole cell lysates.



Immunohistochemical analysis of ACOT4 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with

sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ACOT4 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Anti-ACOT4 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human ACOT4. The exact sequence is proprietary.