

**Anti-PRAS40 Antibody**  
Rabbit polyclonal antibody to PRAS40  
Catalog # AP60425

**Specification**

---

**Anti-PRAS40 Antibody - Product Information**

Application	WB, IH
Primary Accession	<a href="#">O96B36</a>
Other Accession	<a href="#">O9D1F4</a>
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27383

**Anti-PRAS40 Antibody - Additional Information**

**Gene ID** 84335

**Other Names**

PRAS40; Proline-rich AKT1 substrate 1; 40 kDa proline-rich AKT substrate

**Target/Specificity**

Recognizes endogenous levels of PRAS40 protein.

**Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

IH~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-PRAS40 Antibody - Protein Information**

**Name** AKT1S1 {ECO:0000312|EMBL:AAH16043.1}

**Function**

Negative regulator of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:<a href="http://www.uniprot.org/citations/17277771" target="\_blank">17277771</a>, PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/17510057" target="\_blank">17510057</a>, PubMed:<a href="http://www.uniprot.org/citations/29236692" target="\_blank">29236692</a>). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1

activity by blocking the MTOR substrate- recruitment site (PubMed:<a href="http://www.uniprot.org/citations/29236692" target="\_blank">29236692</a>). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/18372248" target="\_blank">18372248</a>). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed:<a href="http://www.uniprot.org/citations/16174443" target="\_blank">16174443</a>, PubMed:<a href="http://www.uniprot.org/citations/18372248" target="\_blank">18372248</a>). May also play a role in nerve growth factor-mediated neuroprotection (By similarity).

#### Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D1F4}. Note=Found in the cytosolic fraction of the brain. {ECO:0000250|UniProtKB:Q9D1F4}

#### Tissue Location


Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)

### Anti-PRAS40 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-PRAS40 Antibody - Images



Western blot analysis of PRAS40 expression in HepG2 (A), EC9706 (B), rat kidney (C) whole cell lysates.



Immunohistochemical analysis of PRAS40 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-PRAS40 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PRAS40. The exact sequence is proprietary.