

**Anti-GRB2 Antibody**  
Rabbit polyclonal antibody to GRB2  
Catalog # AP60465**Specification**

---

**Anti-GRB2 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P62993</a>
Other Accession	<a href="#">Q60631</a>
Reactivity	Human, Mouse, Rat, Zebrafish, Monkey, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25206

**Anti-GRB2 Antibody - Additional Information****Gene ID** 2885**Other Names**

ASH; Growth factor receptor-bound protein 2; Adapter protein GRB2; Protein Ash; SH2/SH3 adapter GRB2

**Target/Specificity**

Recognizes endogenous levels of GRB2 protein.

**Dilution**

WB~~WB (1/500 - 1/1000)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-GRB2 Antibody - Protein Information****Name** GRB2**Synonyms** ASH**Function**Non-enzymatic adapter protein that plays a pivotal role in precisely regulated signaling cascades from cell surface receptors to cellular responses, including signaling transduction and gene expression (PubMed: [11726515](http://www.uniprot.org/citations/11726515), PubMed: [37626338](http://www.uniprot.org/citations/37626338)). Thus, participates in many biological processes including

regulation of innate and adaptive immunity, autophagy, DNA repair or necroptosis (PubMed:<a href="http://www.uniprot.org/citations/35831301" target="\_blank">35831301</a>, PubMed:<a href="http://www.uniprot.org/citations/37626338" target="\_blank">37626338</a>, PubMed:<a href="http://www.uniprot.org/citations/38182563" target="\_blank">38182563</a>). Controls signaling complexes at the T-cell antigen receptor to facilitate the activation, differentiation, and function of T-cells (PubMed:<a href="http://www.uniprot.org/citations/36864087" target="\_blank">36864087</a>, PubMed:<a href="http://www.uniprot.org/citations/9489702" target="\_blank">9489702</a>). Mechanistically, engagement of the TCR leads to phosphorylation of the adapter protein LAT, which serves as docking site for GRB2 (PubMed:<a href="http://www.uniprot.org/citations/9489702" target="\_blank">9489702</a>). In turn, GRB2 establishes a connection with SOS1 that acts as a guanine nucleotide exchange factor and serves as a critical regulator of KRAS/RAF1 leading to MAPKs translocation to the nucleus and activation (PubMed:<a href="http://www.uniprot.org/citations/12171928" target="\_blank">12171928</a>, PubMed:<a href="http://www.uniprot.org/citations/25870599" target="\_blank">25870599</a>). Functions also a role in B-cell activation by amplifying Ca(2+) mobilization and activation of the ERK MAP kinase pathway upon recruitment to the phosphorylated B-cell antigen receptor (BCR) (PubMed:<a href="http://www.uniprot.org/citations/25413232" target="\_blank">25413232</a>, PubMed:<a href="http://www.uniprot.org/citations/29523808" target="\_blank">29523808</a>). Plays a role in switching between autophagy and programmed necrosis upstream of EGFR by interacting with components of necrosomes including RIPK1 and with autophagy regulators SQSTM1 and BECN1 (PubMed:<a href="http://www.uniprot.org/citations/35831301" target="\_blank">35831301</a>, PubMed:<a href="http://www.uniprot.org/citations/38182563" target="\_blank">38182563</a>). Regulates miRNA biogenesis by forming a functional ternary complex with AGO2 and DICER1 (PubMed:<a href="http://www.uniprot.org/citations/37328606" target="\_blank">37328606</a>). Functions in the replication stress response by protecting DNA at stalled replication forks from MRE11-mediated degradation. Mechanistically, inhibits RAD51 ATPase activity to stabilize RAD51 on stalled replication forks (PubMed:<a href="http://www.uniprot.org/citations/38459011" target="\_blank">38459011</a>). Additionally, directly recruits and later releases MRE11 at DNA damage sites during the homology-directed repair (HDR) process (PubMed:<a href="http://www.uniprot.org/citations/34348893" target="\_blank">34348893</a>).

#### Cellular Location

Nucleus. Cytoplasm. Endosome. Golgi apparatus {ECO:0000250|UniProtKB:Q60631}


#### Anti-GRB2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-GRB2 Antibody - Images





Western blot analysis of GRB2 expression in HEK293T (A), HeLa (B), H1688 (C), mouse lung (D), mouse muscle (E), rat lung (F), rat muscle (G) whole cell lysates.

#### **Anti-GRB2 Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GRB2. The exact sequence is proprietary.