

**Anti-KIF2C (pS95) Antibody**  
Rabbit polyclonal antibody to KIF2C (pS95)  
Catalog # AP60580

**Specification**

---

**Anti-KIF2C (pS95) Antibody - Product Information**

Application	<b>WB, IF</b>
Primary Accession	<a href="#">O99661</a>
Reactivity	<b>Human, Monkey</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>81313</b>

**Anti-KIF2C (pS95) Antibody - Additional Information**

**Gene ID** 11004

**Other Names**

KNSL6; Kinesin-like protein KIF2C; Kinesin-like protein 6; Mitotic centromere-associated kinesin; MCAK

**Target/Specificity**

Recognizes endogenous levels of KIF2C (pS95) protein.

**Dilution**

WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)  
IF~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-KIF2C (pS95) Antibody - Protein Information**

**Name** KIF2C

**Synonyms** KNSL6

**Function**

In complex with KIF18B, constitutes the major microtubule plus-end depolymerizing activity in mitotic cells (PubMed: [21820309](http://www.uniprot.org/citations/21820309)). Regulates the turnover of microtubules at the kinetochore and functions in chromosome segregation during mitosis (PubMed: [19060894](http://www.uniprot.org/citations/19060894)). Plays a role in chromosome congression and is required for the lateral to end- on conversion of the

chromosome-microtubule attachment (PubMed:<a href="http://www.uniprot.org/citations/23891108" target="\_blank">23891108</a>).

#### Cellular Location

Cytoplasm, cytoskeleton. Nucleus {ECO:0000250|UniProtKB:P70096} Chromosome, centromere. Chromosome, centromere, kinetochore. Note=Associates with the microtubule network at the growing distal tip (the plus-end) of microtubules, probably through interaction with MTUS2/TIP150 and MAPRE1 (By similarity). Association with microtubule plus ends is also mediated by interaction with KIF18B. Centromeric localization requires the presence of BUB1 and SGO2. {ECO:0000250|UniProtKB:P70096, ECO:0000269|PubMed:17485487, ECO:0000269|PubMed:21820309}

#### Tissue Location


Expressed at high levels in thymus and testis, at low levels in small intestine, the mucosal lining of colon, and placenta, and at very low levels in spleen and ovary; expression is not detected in prostate, peripheral blood Leukocytes, heart, brain, lung, liver, skeletal muscle, kidney or pancreas. Isoform 2 is testis- specific.

### Anti-KIF2C (pS95) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-KIF2C (pS95) Antibody - Images



Western blot analysis of KIF2C (pS95) expression in H1688 (A), H446 (B) whole cell lysates.



Immunofluorescent analysis of KIF2C (pS95) staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

#### **Anti-KIF2C (pS95) Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human KIF2C. The exact sequence is proprietary.