

Anti-PKC beta (pS661) Antibody

Rabbit polyclonal antibody to PKC beta (pS661) Catalog # AP60614

### Specification

# Anti-PKC beta (pS661) Antibody - Product Information

Application Primary Accession Other Accession Reactivity

Host Clonality Calculated MW WB <u>P05771</u> <u>P68404</u> Human, Mouse, Rat, Rabbit, Monkey, Pig, Bovine Rabbit Polyclonal 76869

## Anti-PKC beta (pS661) Antibody - Additional Information

Gene ID 5579

Other Names PKCB; PRKCB1; Protein kinase C beta type; PKC-B; PKC-beta

Target/Specificity Recognizes endogenous levels of PKC beta (pS661) protein.

Dilution WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200)

**Format** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

#### Anti-PKC beta (pS661) Antibody - Protein Information

Name PRKCB

Synonyms PKCB, PRKCB1

#### Function

Calcium-activated, phospholipid- and diacylglycerol (DAG)- dependent serine/threonine-protein kinase involved in various cellular processes such as regulation of the B-cell receptor (BCR) signalosome, oxidative stress-induced apoptosis, androgen receptor-dependent transcription regulation, insulin signaling and endothelial cells proliferation. Plays a key role in B-cell activation by regulating BCR- induced NF-kappa-B activation. Mediates the activation of the canonical NF-kappa-B pathway (NFKB1) by direct phosphorylation of CARD11/CARMA1 at 'Ser-559', 'Ser-644'



and 'Ser-652'. Phosphorylation induces CARD11/CARMA1 association with lipid rafts and recruitment of the BCL10-MALT1 complex as well as MAP3K7/TAK1, which then activates IKK complex, resulting in nuclear translocation and activation of NFKB1. Plays a direct role in the negative feedback regulation of the BCR signaling, by down-modulating BTK function via direct phosphorylation of BTK at 'Ser-180', which results in the alteration of BTK plasma membrane localization and in turn inhibition of BTK activity (PubMed:<a

href="http://www.uniprot.org/citations/11598012" target=" blank">11598012</a>). Involved in apoptosis following oxidative damage: in case of oxidative conditions, specifically phosphorylates 'Ser-36' of isoform p66Shc of SHC1, leading to mitochondrial accumulation of p66Shc, where p66Shc acts as a reactive oxygen species producer. Acts as a coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and specifically mediating phosphorylation of 'Thr-6' of histone H3 (H3T6ph), a specific tag for epigenetic transcriptional activation that prevents demethylation of histone H3 'Lys-4' (H3K4me) by LSD1/KDM1A (PubMed:<a href="http://www.uniprot.org/citations/20228790" target=" blank">20228790</a>). In insulin signaling, may function downstream of IRS1 in muscle cells and mediate insulin-dependent DNA synthesis through the RAF1-MAPK/ERK signaling cascade. Participates in the regulation of glucose transport in adjpocytes by negatively modulating the insulin-stimulated translocation of the glucose transporter SLC2A4/GLUT4. Phosphorylates SLC2A1/GLUT1, promoting glucose uptake by SLC2A1/GLUT1 (PubMed:<a href="http://www.uniprot.org/citations/25982116" target=" blank">25982116</a>). Under high glucose in pancreatic beta-cells, is probably involved in the inhibition of the insulin gene transcription, via regulation of MYC expression. In endothelial cells, activation of PRKCB induces increased phosphorylation of RB1, increased VEGFA-induced cell proliferation, and inhibits PI3K/AKT-dependent nitric oxide synthase (NOS3/eNOS) regulation by insulin, which causes endothelial dysfunction. Also involved in triglyceride homeostasis (By similarity). Phosphorylates ATF2 which promotes cooperation between ATF2 and JUN, activating transcription (PubMed: <a href="http://www.uniprot.org/citations/19176525" target=" blank">19176525</a>). Phosphorylates KLHL3 in response to angiotensin II signaling, decreasing the interaction between

KLHL3 and WNK4 (PubMed:<a href="http://www.uniprot.org/citations/25313067" target="\_blank">25313067</a>). Phosphorylates and activates LRRK1, which phosphorylates RAB proteins involved in intracellular trafficking (PubMed:<a href="http://www.uniprot.org/citations/36040231" target=" blank">36040231</a>).

**Cellular Location** 

Cytoplasm. Nucleus. Membrane; Peripheral membrane protein

# Anti-PKC beta (pS661) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-PKC beta (pS661) Antibody - Images





Western blot analysis of PKC beta (pS661) expression in mouse brain (A) whole cell lysates.



Immunohistochemical analysis of PKC beta (pS661) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# Anti-PKC beta (pS661) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PKC beta. The exact sequence is proprietary.